For Research Use Only

## CoraLite® Plus 750 Anti-Human TLR3/CD283 Rabbit Recombinant Antibody



Catalog Number: CL750-98016-2

**Basic Information** 

Catalog Number:

CL750-98016-2

Size: 100tests, 5 ul/test

Source: Rabbit Isotype:

Immunogen Catalog Number:

GenBank Accession Number: BC094737

GeneID (NCBI):

UNIPROT ID: O15455 Full Name:

7098

toll-like receptor 3

Calculated MW:
904 aa, 104 kDa

Purification Method:

Protein A purification CloneNo.:

Excitation/Emission maxima

wavelengths: 755 nm / 780 nm

230445G9

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity:

human

**Background Information** 

TLR3 belongs to the Toll-like receptor family which is important in the innate immune response to pathogens. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. TLR3 is a nucleotide-sensing TLR that is activated by double-stranded RNA. Upon recognition of dsRNA, TLR3 transmits signals via the adaptor protein TICAM-1/TRIF, leading to the induction of type I IFN, cytokine/chemokine production, and dendritic cell (DC) maturation (PMID: 18262679).

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide and 0.5% BSA.

## **Selected Validation Data**





1x10^6 human monocyte-derived immature dendritic cells were intracellularly stained with 5 ul CoraLite® Plus 750 Anti-Human TLR3/CD283 Rabbit Recombinant Antibody (CL750-98016-2, Clone: 230445G9) (right), or Isotype Control (left). Cells were co-stained with CoraLite® Plus 488 Anti-Human CD11c. Cells were incubated with FC Receptor Block prior to fixing and staining. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1x10^6 human monocyte-derived immature dendritic cells were intracellularly stained with 5 ul CoraLite® Plus 750 Anti-Human TLR3/CD283 Rabbit Recombinant Antibody (CL750-98016-2, Clone: 230445G9) (red), or Isotype Control (blue). Cells were incubated with FC Receptor Block prior to fixing and staining. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).