For Research Use Only

CoraLite® Plus 750 Anti-Mouse ICAM-1/CD54 Rabbit Recombinant Antibody

Catalog Number: CL750-98049



Purification Method:

Protein A purification

755 nm / 780 nm

Basic Information

Applications

Catalog Number: GenBank Accession Number: CL750-98049 NM_010493

GeneID (NCBI): CloneNo.: Concentration: 100ug, 500 $\,\mu$ g/ml 15894 240249D10

UNIPROT ID: Source: Recommended Dilutions:

Rabbit P13597 FC: 0.25 ug per 10^6 cells in a 100 μ l suspension

Full Name: Isotype:

intercellular adhesion molecule 1 Excitation/Emission maxima wavelengths:

Calculated MW: Immunogen Catalog Number: EG0482

59 kDa

Positive Controls: **Tested Applications:**

FC: mouse splenocytes, Species Specificity:

Background Information

ICAM-1 (CD54) is a 90-kDa transmembrane glycoprotein of the immunoglobulin superfamily and is critical for the firm attachment and transmigration of leukocytes out of blood vessels and into tissues (PMID: 19307690). ICAM-1 is expressed by several cell types, typically on endothelial cells and cells of the immune system, and its expression can be up-regulated by various stimuli, including TNF- α , INF- γ , IL-1 and thrombin (PMID: 3086451; 9694714; 15979056). It is a ligand for LFA-1 and Mac-1, serves as a receptor for rhinovirus, and is one of several receptors used by Plasmodium falciparum (PMID: 2566624; 2538244; 2475784).

Storage

mouse

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide, pH7.3

Selected Validation Data





1x10^6 mouse splenocytes were surface stained with CoraLite® Plus 488 Anti-Mouse CD3, and 0.25 ug CoraLite® Plus 750 Anti-Mouse ICAM-1/CD54 Rabbit Recombinant Antibody (CL750-98049, Clone: 240249D10) or Isotype Control. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 mouse splenocytes were surface stained with 0.25 ug CoraLite® Plus 750 Anti-Mouse ICAM-1/CD54 Rabbit Recombinant Antibody (CL750-98049, Clone: 240249D10) (red) or 0.25 ug Isotype Control (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.