

Catalog Number: CM06238

## 产品信息

**Catalog Number:**  
CM06238

**CAS号:**  
517-88-4


**分子式:**  
C<sub>16</sub>H<sub>16</sub>O<sub>5</sub>

**主要靶点:**  
PI3K|Apoptosis

**主要通路:**  
PI3K/Akt/mTOR信号通路|凋亡

**分子量:**  
288.3

**溶解度:**  
DMSO:50 mg/mL (173.43 mM)



## 体外活性

3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay was performed for cell viability and adhesion ability of MGC-803 cells. Cell scratch repair experiments were conducted for the determination of migration ability while transwell assay for cell invasion ability. Western blot analysis and real-time polymerase chain reaction assay were used for the detection of protein and mRNA expressions. Fifty per cent inhibitory concentration of Shikonin on MGC-803 cells was 1.854 μm. Shikonin (1 μm) inhibited significantly the adhesion, invasion and migratory ability of MGC-803 cells. Interestingly, Shikonin in the presence or absence of anti-Toll-like receptor 2 (TLR2) antibody (2 μg) and nuclear factor-kappa B (NF-κB) inhibitor MG-132 (10 μm) could decrease these ability of MGC-803 cells markedly, as well as the expression levels of matrix metalloproteinases (MMP)-2, MMP-7, TLR2 and p65 NF-κB. In addition, the co-incubation of Shikonin and anti-TLR2/MG-132 has a significant stronger activity than anti-TLR2 or MG-132 alone.

## 描述

(-)-Alkannin (Shikonin) is a natural red naphthoquinone pigment, has antimicrobial, anti-tumor, and anti-inflammatory effects; a purified shikonin preparation is widely used for the production of medicinals, cosmetics, and some food products; shikonin also enters into the antiinflammatory ointment and cream compositions used for the treatment of burns. It can suppress the cell viability, adhesion, invasion and migratory ability of MGC-803 cells through TLR2- or NF-κB-mediated pathway.

## 储存

Powder: -20°C for 3 years | In solvent: -80°C for 2 years