## For Research Use Only

## PerCP-Cyanine 5.5 Anti-Human CD8 (UCHT4) Mouse IgG2a Recombinant Antibody



Catalog Number: CPY5-65559

**Basic Information** 

Catalog Number: CPY5-65559

Size:

100tests, 5 µl/test

Source: Mouse Isotype:

IgG2a

GenBank Accession Number: BC025715

GeneID (NCBI):

ENSEMBL Gene ID: ENSG00000153563

**UNIPROT ID:** P01732 Full Name: CD8a molecule

Calculated MW: 235 aa, 26 kDa

**Purification Method:** 

Protein A purification

CloneNo.: UCHT4

Excitation/Emission maxima

wavelengths: 482 nm / 690 nm

**Applications** 

**Tested Applications:** 

Species Specificity:

human

## **Background Information**


CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8  $\alpha$  or as a heterodimer of CD8  $\alpha$  and CD8  $\beta$  (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8  $\alpha$   $\beta$  heterodimer while CD8  $\alpha$   $\alpha$  homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR)  $\gamma$   $\delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

## Selected Validation Data



1x10^6 human PBMCs were surface stained with 5 ul PerCP-Cyanine5.5 CD8 Recombinant Antibody (CPY5-65559, Clone: UCHT4) or unstained. Cells were co-stained with CoraLite® Plus 647 Anti-Human CD3 (UCHT1) (CL647-65151, Clone: UCHT1). Cells were not fixed.