

For Research Use Only

FcZero-rAb™ PerCP-Cyanine5.5 Anti-Human CD8 (SK1) Rabbit IgG Recombinant Antibody

Catalog Number:CPY5-FcA65630



Basic Information

Catalog Number:	CPY5-FcA65630	GenBank Accession Number:	BC025715	Purification Method:
Concentration:	100 tests, 5 μ l/test	GenID (NCBI):	925	Protein A purification
Source:	Rabbit	ENSEMBL Gene ID:	ENSG00000153563	Clone No.:
Isotype:	IgG	UNIPROT ID:	P01732	SK1
		Full Name:	CD8a molecule	Recommended Dilutions:
		Calculated MW:	235 aa, 26 kDa	FC: 5 μ l per 10^6 cells in a 100 μ l suspension
				Excitation/Emission maxima wavelengths: 482 nm / 690nm

Applications

Tested Applications:	Positive Controls:
FC	FC : human PBMCs,
Species Specificity:	human, monkey, non-human primates

Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α α homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γ δ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).

Storage

Storage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human PBMCs were surface stained with FcZero-rAb™ CoraLite® Plus 488 Anti-Human CD3 (OKT3), and 5 μ l PerCP-Cyanine5.5 Anti-Human CD8 (SK1) Rabbit IgG RecAb (CPY5-FcA65630, Clone:SK1) or FcZero-rAb™ PerCP-Cyanine5.5 Rabbit IgG Isotype Control Recombinant Antibody (CPY5-FcA98136, Clone: 240953C9). Cells were incubated with MonoZero™ Monocytes blocking Reagent (PF00020) prior to staining. Cells were not fixed. Lymphocytes were

1x10⁶ human PBMCs were surface stained with 5 μ l PerCP-Cyanine5.5 Anti-Human CD8 (SK1) Rabbit IgG RecAb (CPY5-FcA65630, Clone:SK1) (red) or FcZero-rAb™ PerCP-Cyanine5.5 Rabbit IgG Isotype Control Recombinant Antibody (CPY5-FcA98136, Clone: 240953C9) (blue). Cells were incubated with MonoZero™ Monocytes blocking Reagent (PF00020) prior to staining. Cells were not fixed. Lymphocytes were gated.