## For Research Use Only

## Cardinal Red™-conjugated Cyclin E2 Polyclonal antibody




Catalog Number:CR-11935

| Basic Information      | Catalog Number:<br>CR-11935  | GenBank Accession Number:<br>BC020729                                      | Purification Method:<br>Antigen affinity purification         |  |  |  |
|------------------------|--|--|---|--|--|--|
|                        | Size:<br>1000 µg/ml<br>Source:<br>Rabbit<br>Isotype:<br>IgG<br>Immunogen Catalog Number:<br>AG2532   | GeneID (NCBI):<br>9134<br>UNIPROT ID:<br>096020<br>Full Name:<br>cyclin E2 | Excitation/Emission maxima<br>wavelengths:<br>592 nm / 611 nm |  |  |  |
|                        |  |  |   | Calculated MW:<br>374 aa, 44 kDa                           |  |  |
|                        |  |  |   | Observed MW:<br>44 kDa                                     |  |  |
|                        |  | Applications   |   | Tested Applications:<br>FC (Intra)<br>Species Specificity: |  |  |
|                        |  |  |   | human  |  |  |
| Background Information | Cyclin E2 (CCNE2) belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. Cyclins function as regulators of cyclindependent kinases (CDKs). Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of cell cycle events. CCNE2 forms a complex with and functions as a regulatory subunit of CDK2 and has been shown to specifically interact with CIP/KIP family of CDK inhibitors. CCNE2 plays a role in cell cycle G1/S transition and its expression peaks at the G1-S phase. Whereas cyclin E1 is expressed in most proliferating normal and tumor cells, cyclin E2 levels are low or undetectable in nontransformed cells, and are elevated in tumor-derived cells. |  |   |  |  |  |
| Storage                | Storage:<br>Store at -20°C. Avoid exposure to l<br>Storage Buffer:<br>PBS with 50% Glycerol, 0.05% Pro<br>Aliquoting is unnecessary for -20°   |  | nt.   |  |  |  |

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.2 ug Cardinal Red™ Anti-Human Cyclin E2 (CR-11935) (red), or 0.2 ug isotype control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).