

For Research Use Only

Recombinant Human TNFR2/CD120b protein (Myc Tag, His Tag)



Catalog Number: Eg0086

Basic Information

Species:
Human

EC50:
1-5 ng/mL

Purity:
>90 %, SDS-PAGE

Tag:
Myc Tag, His Tag

Technical Specifications

Purity:
>90 %, SDS-PAGE

Endotoxin Level:
<0.1 EU/ μ g protein, LAL method

Source:
HEK293-derived Human TNFR2 protein Leu23-Asp257 (Accession# P20333-1) with a Myc tag and a His tag at the C-terminus.

GeneID:
7133

Accession:
P20333-1

Predicted Molecular Mass:
30.4 kDa

SDS-PAGE:
48-58 kDa, reducing (R) conditions

Formulation:
Lyophilized from 0.22 μ m filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

Biological Activity

1. Immobilized Human TNF- α (GST tag) at 2 μ g/mL (100 μ L/well) can bind Human TNFR2 (Myc tag, His tag) with a linear range of 1-5 ng/mL.
2. Measured by its ability to inhibit TNF- α mediated cytotoxicity in the L-929 mouse fibroblast cells in the presence of the metabolic inhibitor actinomycin D. The EC50 for this effect is 50-200 ng/mL in the presence of 0.25 ng/mL recombinant human TNF- α .

Storage and Shipping

Storage:

It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

- Until expiry date, -20°C to -80°C as lyophilized proteins.
- 3 months, -20°C to -80°C under sterile conditions after reconstitution.

Shipping:

The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

Background

Tumor necrosis factor- α (TNFA/TNFSF2) is a multifunctional cytokine that plays a key role in regulating inflammation, immune functions, host defense, and apoptosis. TNFA signals through two distinct cell surface receptors, TNFR1 (TNFRSF1A, CD120a, p55) and TNFR2 (TNFRSF1B, CD120b, p75). TNFR2 is a kind of receptor with high affinity for TNFSF2/TNF- α and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin- α . The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF- α . In contrast to TNFR1, TNFR2 does not have a death domain. TNFR2 only signals for antiapoptotic reactions. However, recent evidence indicates that TNFR2 also signals to induce TRAF2 degradation. Various defects in the TNFR2 pathway, due to polymorphisms in the TNFR2 gene, upregulated expression of TNFR2 and TNFR2 shedding, have been implicated in the pathology of several autoimmune disorders.


References

1. Islam A. et al. (2006). J Biol Chem. 281(10):6860-73.
2. Pan S. et al. (2002). Mol Cell Biol. 22(21):7512-23.
3. Faustman D. et al. (2010). Nat Rev Drug Discov. 9(6):482-93.


Synonyms

TNFRSF1B, CD120b, Etanercept, p75, p75TNFR


Selected Validation Data



Purity of Recombinant Human TNFR2 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) conditions and stained using Coomassie blue.



Immobilized Human TNF- α (GST tag) at 2 μ g/mL (100 μ L/well) can bind Human TNFR2 (Myc tag, His tag) with a linear range of 1-5 ng/mL



Measured by its ability to inhibit TNF- α mediated cytotoxicity in the L-929 mouse fibroblast cells in the presence of the metabolic inhibitor actinomycin D. The EC₅₀ for this effect is 50-200 ng/mL in the presence of 0.25 ng/mL recombinant human TNF- α .

For technical support and original validation data for this product please contact

T: 027-87531629

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.