For Research Use Only

Recombinant Human CD14 protein (His Tag)



www.ptgcn.com

Catalog Number: Eg0498

Basic Information

Species: Human

Purity: >90 %, SDS-PAGE

Tag: His Tag

Technical Specifications

Purity: >90 %, SDS-PAGE

Endotoxin Level:

<0.1 EU/ µ g protein, LAL method

HEK293-derived Human CD14 protein Thr20-Met344 (Accession# P08571) with a His tag at the C-terminus.

GeneID:

Accession: P08571

Predicted Molecular Mass:

39.1 kDa

SDS-PAGE:

45-60 kDa, reducing (R) conditions

Lyophilized from 0.22 $\,\mu$ m filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

Biological Activity

Not tested

Storage and Shipping

It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Until expiry date, -20°C to -80°C as lyophilized proteins.

3 months, -20℃ to -80℃ under sterile conditions after reconstitution.

The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

Background

CD14 is a 50-55 kDa glycosylphosphatidylinositol-anchored glycoprotein preferentially expressed on monocytes and macrophages, and at lower levels on granulocytes. CD14 can also exist as a soluble protein. CD14 acts as a co-receptor for bacterial liposaccharides (LPS). It plays a major role in the inflammatory response of monocytes to LPS.


References

1. A Haziot, et al. (1988) J Immunol. 141(2):547-52. 2. D L Simmons, et al. (1989) Blood. 73(1):284-9. 3. S D Wright, et al. (1990) Science. 249(4975):1431-3. 4. A Haziot, et al. (1993) J Immunol. 150(12):5556-65.

Synonyms

CD14, CD14 molecule, Monocyte differentiation antigen CD14, membrane-bound form, Monocyte differentiation antigen CD14, urinary form, Myeloid cell-specific leucine-rich glycoprotein

Selected Validation Data



Purity of Recombinant Human CD14 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.