

Recombinant Mouse E-Cadherin protein (His Tag)

Catalog Number: Eg0975

Basic Information

Species:
Mouse

Purity:
>90 %, SDS-PAGE

Tag:
His Tag

Technical Specifications

Purity:

>90 %, SDS-PAGE

Endotoxin Level:

<0.1 EU/ µg protein, LAL method

Source:

HEK293-derived Mouse E-Cadherin protein Asp157-Val709 (Accession# P09803) with a His tag at the C-terminus.

GeneID:

12550

Accession:

P09803

Predicted Molecular Mass:

61.9 kDa

SDS-PAGE:

65-75 kDa, reducing (R) conditions

Formulation:

Lyophilized from 0.22 µm filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

Biological Activity

Not tested

Storage and Shipping

Storage:

It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

- Until expiry date, -20°C to -80°C as lyophilized proteins.
- 3 months, -20°C to -80°C under sterile conditions after reconstitution.

Shipping:

The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

Background

E-cadherin (epithelial cadherin), also known as CDH1 (cadherin 1) or CAM 120/80, is a classical member of the cadherin superfamily which also include N-, P-, R-, and B-cadherins. E-cadherin is expressed on the cell surface in most epithelial tissues. The extracellular region of E-cadherin establishes calcium-dependent homophilic trans binding, providing specific interaction with adjacent cells, while the cytoplasmic domain is connected to the actin cytoskeleton through the interaction with p120-, α-, β-, and γ-catenin (plakoglobin). E-cadherin is important in the maintenance of the epithelial integrity, and is involved in mechanisms regulating proliferation, differentiation, and survival of epithelial cell. E-cadherin may also play a role in tumorigenesis. It is considered to be an invasion suppressor protein and its loss is an indicator of high tumor aggressiveness. E-cadherin is sensitive to trypsin digestion in the absence of Ca²⁺.


References

1. Braga V. et al. (2016) Curr Opin Cell Biol. 42:138-21.
2. Ishiyama N. et al. (2017) Cold Spring Harb Perspect Biol. 9(5):a028738.
3. Dongre A. et al. (2019) Nat Rev Mol Cell Biol. 20(2):69-84.
4. Yang, S. et al. (2013) Biomed res int. 2013:608728.
5. Harrison OJ. et al. (2010) Nat Struct Mol Biol. 17(3):348-357.
6. Biswas KH. et al. (2017) Exp Cell Res. 358(1):14-19.
7. Buckley CD. et al. (2014) Science. 346(6209):1254211.

Synonyms

E-cadherin, cadherin 1, Cdh1, Ecad, L CAM

Selected Validation Data



Purity of Recombinant Mouse E-Cadherin was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.

For technical support and original validation data for this product please contact

T: 027-87531629

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.