## For Research Use Only

## Recombinant Human Kininogen-1 protein (His Tag)



Catalog Number: Eg1162

**Basic Information** 

Species: Human

Purity: >90 %, SDS-PAGE

Tag: His Tag

**Technical Specifications** 

Purity: >90 %, SDS-PAGE

**Endotoxin Level:** 

<0.1 EU/  $\mu$  g protein, LAL method

HEK293-derived Human Kininogen-1 protein Gln19-Ser644 (Accession# P01042-1) with a His tag at the Cterminus.

GeneID:

3827

**Accession:** 

P01042-1

**Predicted Molecular Mass:** 

71.5 kDa

**SDS-PAGE**:

95-110 kDa, reducing (R) conditions

Formulation:
Lyophilized from 0.22 µm filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

**Biological Activity** 

Not tested

Storage and Shipping

Storage:

It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Until expiry date, -20% to -80% as lyophilized proteins. 3 months, -20% to -80% under sterile conditions after reconstitution.

Shipping:
The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

Background

Kininogens are inhibitors of thiol proteases. Kininogen 1 plays important role in Kinin-kallikrein system. This gene is translated into High-molecular weight kininogen (HMWK) and low-molecular weight kininogen (LMWK) after alternative splicing. HMWK is produced by the liver together with prekallikrein. It acts mainly as a cofactor on coagulation and inflammation, and has no intrinsic catalytic activity. LMWK is produced locally by numerous tissues, and secreted together with tissue kallikrein.


References

1. Cheng Z. et al. (2018) J Exp Clin Cancer Res. 37(1):180.

**Synonyms** 

KNG1, Alpha-2-thiol proteinase inhibitor, BDK, Bradykinin, Fitzgerald factor

## **Selected Validation Data**



Purity of Recombinant Human Kininogen-1 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.