For Research Use Only

Recombinant Human NKp46/NCR1 protein (rFc Tag)



Catalog Number: Eg1364

Basic Information

Species: Human

Purity: >90 %, SDS-PAGE

Tag: rFc Tag

Technical Specifications

Purity: >90 %, SDS-PAGE

Endotoxin Level:

<0.1 EU/ µ g protein, LAL method

HEK293-derived Human NKp46 protein Gln22-Arg258 (Accession# EAW72309.1) with a rabbit IgG Fc tag at the Cterminus.

GeneID:

9437

Accession: EAW72309.1

Predicted Molecular Mass:

53.1 kDa

SDS-PAGE:

55-70 kDa, reducing (R) conditions

Formulation:
Lyophilized from 0.22 μm filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

Biological Activity

Not tested

Storage and Shipping

Storage:

It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Until expiry date, -20% to -80% as lyophilized proteins. 3 months, -20% to -80% under sterile conditions after reconstitution.

Shipping:
The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

Background

NKp46, also known as NCR1 or CD335, is an Ig-like superfamily cell surface receptor that is highly conserved in mammals. NKp46 is a type I transmembrane glycoprotein consisting of two extracellular Ig-like domains, a transmembrane domain, and an intracellular tail. It is expressed on NK cells, rare T-cell subsets and a mucosal population of NKp46+ innate lymphoid cells. NKp46 is the major triggering receptor involved in the natural cytotoxicity.

References


- 1. Narni-Mancinelli E. et al. (2011). Proc Natl Acad Sci U S A. 108(45):18324-9. 2. Pessino A. et al. (1998). J Exp Med. 188(5):953-60. 3. Sivori S. et al. (1999). Eur J Immunol. 29(5):1656-66. 4. Biassoni R. et al. (1999). Eur J Immunol. 29(3):1014-20.

- 5. Westgaard IH. et al. (2004). J Leukoc Biol. 76(6):1200-6.

Synonyms

CD335, hNKp46, LY94, Natural killer cell p46-related protein, NCR 1

Selected Validation Data



Purity of Recombinant Human NKp46 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.