## For Research Use Only Recombinant Human CD63 protein (rFc Tag)



www.ptgcn.com

## Catalog Number: Eg2064

| Basic Information          | <mark>Species:</mark><br>Human   | Purity:<br>>90 %, SDS-PAGE  | Tag:<br>rFc Tag                   |
|----------------------------|--|---|-----------------------------------|
| Technical Specifications   | Purity:<br>>90 %, SDS-PAGE   |   |                                   |
|                            | <mark>Endotoxin Level:</mark><br><0.1 EU/μg protein, LAL method  |   |                                   |
|                            | Source:<br>HEK293-derived Human CD63 protein Ala103-Val203 (Accession# P08962-1) with a rabbit IgG Fc tag at the C-<br>terminus.   |   |                                   |
|                            | GenelD:<br>967   |   |                                   |
|                            | Accession:<br>P08962-1   |   |                                   |
|                            | Predicted Molecular Mass:<br>37.8 kDa  |   |                                   |
|                            | SDS-PAGE:<br>42-58 kDa, reducing (R) conditions  |   |                                   |
|                            | <b>Formulation:</b><br>Lyophilized from 0.22 μ m filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as<br>protectants before lyophilization.  |   |                                   |
| <b>Biological Activity</b> | Not tested   |   |                                   |
| Storage and Shipping       | <b>Storage:</b><br>It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.   |   |                                   |
|                            | <ul> <li>Until expiry date, -20℃ to -80℃ as lyophilized proteins.</li> <li>3 months, -20℃ to -80℃ under sterile conditions after reconstitution.</li> </ul>  |   |                                   |
|                            | <b>Shipping:</b><br>The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended<br>temperature.   |   |                                   |
| Reconstitution             | Briefly centrifuge the   | ube before opening. Reconstitute at 0.1-0.5   | mg/mL in sterile water.           |
| Background                 | CD63 is a 30-60 kDa lysosomal membrane protein that belongs to the tetraspanin family. This protein plays<br>many important roles in immuno-physiological functions. It mediates signal transduction events that play a<br>role in the regulation of cell development, activation, growth, and motility. CD63 is expressed on activated<br>platelets, thus it may function as a blood platelet activation marker. CD63 is a lysosomal membrane<br>glycoprotein that is translocated to plasma membrane after platelet activation. The CD63 tetraspanin is<br>highly expressed in the early stages of melanoma and decreases in advanced lesions, suggesting it as a<br>possible suppressor of tumor progression. Deficiency of this protein is associated with Hermansky-Pudlak<br>syndrome. |   |                                   |
| References                 | 2. Guillaume van Niel, e<br>3. Maaike S Pols, Judith<br>4. D O Azorsa, et al. (199   | (2006) EMBO J. Sep 6;25(17):3934-42.<br>t al. (2011) Dev Cell. Oct 18;21(4):708-21.<br>Klumperman. (2009) Exp Cell Res. May 15;31!<br>91) Blood. Jul 15;78(2):280-4.<br>93) J Clin Invest. Apr;91(4):1775-82.Maaike S |                                   |
| Synonyms                   | CD 63, CD63 antigen, Li<br>protein 1   | mp1, Lysosomal-associated membrane prot   | ein 3, Lysosome integral membrane |

## Selected Validation Data



Purity of Recombinant Human CD63 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.