

# Recombinant Human IL-10RA protein (rFc Tag)

Catalog Number: Eg2314

## Basic Information

**Species:**  
Human

**Purity:**  
>90 %, SDS-PAGE

**Tag:**  
rFc Tag

## Technical Specifications

**Purity:**

>90 %, SDS-PAGE

**Endotoxin Level:**

<0.1 EU/ µg protein, LAL method

**Source:**

HEK293-derived Human IL-10RA protein His22-Asn235 (Accession# Q13651) with a rabbit IgG Fc tag at the C-terminus.

**GeneID:**

3587

**Accession:**

Q13651

**Predicted Molecular Mass:**

50.7 kDa

**SDS-PAGE:**

60-85 kDa, reducing (R) conditions

**Formulation:**

Lyophilized from 0.22 µm filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

## Biological Activity

Not tested

## Storage and Shipping

**Storage:**

It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

- Until expiry date, -20°C to -80°C as lyophilized proteins.
- 3 months, -20°C to -80°C under sterile conditions after reconstitution.

**Shipping:**

The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

## Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

## Background

Interleukin-10 (IL-10) is an anti-inflammatory cytokine that plays a critical role in maintaining immune system balance. IL-10 signals through a receptor complex consisting of two molecules of the IL-10R alpha-chain (IL-10RA) and two molecules of the accessory IL-10R beta-chain (IL-10RB) (PMID: 22236434). IL-10RA and IL-10RB are members of the type II cytokine receptor family. IL-10RA, also known as IL-10R1 or CD210, is expressed on most hematopoietic cells at a basal level but is upregulated by various cells upon activation (PMID: 24507158). IL-10RA serves as the ligand-binding subunit of the receptor complex. Upon binding of IL-10 to the IL-10R complex, the kinases JAK1 and TYK2 are activated and catalyze the phosphorylation of IL-10RA at specific intracellular tyrosine residues, leading to the recruitment and subsequent phosphorylation of STAT3 (PMID: 11244051; 10433356).


## References

1. Glocker EO. et al. (2011). Ann N Y Acad Sci. 1246:102-107.
2. Shouval DS. et al. (2014). Adv Immunol. 122:177-210.
3. Moore KW. et al. (2001). Annu Rev Immunol. 19:683-765.
4. Donnelly RP. et al. (1999). J Interferon Cytokine Res. 19(6):563-573.

## Synonyms

IL10RA, CD210, HIL 10R, IL 10 receptor subunit alpha, IL 10R subunit 1

## Selected Validation Data



Purity of Recombinant Human IL-10RA was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.

For technical support and original validation data for this product please contact

T: 027-87531629

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.