

Recombinant Human Progranulin/PGRN protein (rFc Tag)

Catalog Number: Eg2954

Basic Information

Species:
Human**Purity:**
>90 %, SDS-PAGE**Tag:**
rFc Tag

Technical Specifications

Purity:

>90 %, SDS-PAGE

Endotoxin Level:<0.1 EU/ μ g protein, LAL method**Source:**

HEK293-derived Human Progranulin protein Thr18-Leu593 (Accession# P28799-1) with a rabbit IgG Fc tag at the C-terminus.

GeneID:

2896

Accession:

P28799-1

Predicted Molecular Mass:

87.8 kDa

SDS-PAGE:

95-105 kDa, reducing (R) conditions

Formulation:Lyophilized from 0.22 μ m filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

Biological Activity

Not tested

Storage and Shipping

Storage:

It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

- Until expiry date, -20°C to -80°C as lyophilized proteins.
- 3 months, -20°C to -80°C under sterile conditions after reconstitution.

Shipping:

The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

Background

GRN, also known as PGRN or PCDGF, is a cysteine-rich protein of 68.5 kDa that is typically secreted into a highly glycosylated 88 kDa form. PGRN is a unique growth factor that plays an important role in cutaneous wound healing. It has an anti-inflammatory effect and promotes cell proliferation. When PCDGF is degraded to several 6-25 kDa fragments, called granulins (GRNs) by neutrophil proteases, a pro-inflammatory reaction occurs. PGRN is widely expressed, particularly in epithelial cells, immune cells, neurons, and chondrocytes. High levels of PGRN expression have been reported in human cancers, and its expression is closely correlated with the development and metastasis of several cancers. The recent discovery that mutations in the gene encoding for pro-granulin (GRN) cause frontotemporal lobar degeneration (FTLD), and other neurodegenerative diseases leading to dementia, has brought renewed interest in progranulin and its functions in the central nervous system.


References

1. Ahmed Z. et al. (2007). J Neuroinflammation. 4:7.

Synonyms

granulin, Epithelin precursor, Glycoprotein 88, Glycoprotein of 88 Kda, Granulin precursor

Selected Validation Data



Purity of Recombinant Human Progranulin was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) conditions and stained using Coomassie blue.

For technical support and original validation data for this product please contact

T: 027-87531629

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.