

For Research Use Only

Recombinant Human CD19 protein (rFc Tag)



Catalog Number: Eg3335

Basic Information

Species:
Human

Purity:
>90 %, SDS-PAGE

Tag:
rFc Tag

Technical Specifications

Purity:

>90 %, SDS-PAGE

Endotoxin Level:

<0.1 EU/ μ g protein, LAL method

Source:

HEK293-derived Human CD19 protein Pro20-Lys291 (Accession# P15391-1) with a rabbit IgG Fc tag at the C-terminus.

GeneID:

930

Accession:

P15391-1

Predicted Molecular Mass:

56.4 kDa

SDS-PAGE:

65-80 kDa, reducing (R) conditions

Formulation:

Lyophilized from 0.22 μ m filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

Biological Activity

Not tested

Storage and Shipping

Storage:

It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

- Until expiry date, -20°C to -80°C as lyophilized proteins.
- 3 months, -20°C to -80°C under sterile conditions after reconstitution.

Shipping:

The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

Background

CD19 is a type I transmembrane glycoprotein belonging to the immunoglobulin superfamily (PMID: 2472450). It is expressed by B cells and follicular dendritic cells. CD19 is up-regulated at the step of B-lineage commitment during the differentiation of the hematopoietic stem cell, it remains on during subsequent stages of differentiation until finally down-regulated during terminal differentiation into plasma cells (PMID: 8528044). CD19 is involved in B cell development, activation and differentiation. It is the dominant component for the signaling complex on B cells that includes CD21 (CR2), CD81 (TAPA-1) and CD225 and acts as a critical co-receptor for BCR signal transduction (PMID: 23210908).


References

1. Tedder, T F, and C M Isaacs. Journal of immunology (Baltimore, Md. : 1950) vol. 143,2 (1989): 712-7.
2. Scheuermann, R H, and E Racila. Leukemia & lymphoma vol. 18,5-6 (1995): 385-97.
3. Wang, Kemeng et al. Experimental hematology & oncology vol. 1,1(2012) 36.

Synonyms

B lymphocyte antigen CD19, B4, B-lymphocyte antigen CD19, B-lymphocyte surface antigen B4, CD19 molecule

Selected Validation Data



Purity of Recombinant Human CD19 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) conditions and stained using Coomassie blue.

For technical support and original validation data for this product please contact

T: 027-87531629

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.