

Recombinant Human NRG2 protein (rFc Tag)

Catalog Number: Eg4172

Basic Information

Species:
Human

Purity:
>90 %, SDS-PAGE

Tag:
rFc Tag

Technical Specifications

Purity:

>90 %, SDS-PAGE

Endotoxin Level:

<0.1 EU/ µg protein, LAL method

Source:

HEK293-derived Human NRG2 protein Cys112-Lys404 (Accession# O14511-1) with a rabbit IgG Fc tag at the C-terminus.

GeneID:

9542

Accession:

O14511-1

Predicted Molecular Mass:

58.9 kDa

SDS-PAGE:

65-85 kDa, reducing (R) conditions

Formulation:

Lyophilized from 0.22 µm filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

Biological Activity

Not tested

Storage and Shipping

Storage:

It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

- Until expiry date, -20°C to -80°C as lyophilized proteins.
- 3 months, -20°C to -80°C under sterile conditions after reconstitution.

Shipping:

The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

Background

NRG2 is a novel member of the neuregulin family of growth and differentiation factors. By interacting with the ERBB family of receptors, this protein induces the growth and differentiation of epithelial, neuronal, glial and other cell types. The gene consists of 12 exons and the genomic structure is similar to that of neuregulin 1, another member of the neuregulin family of ligands. The products of these genes mediate different biological processes by acting at different sites in tissues and eliciting different biological responses in cells. This gene is close to the Charcot-Marie-Tooth demyelinating disease locus, but does not cause the disease.


References

1. Yan L. et al. (2018) Mol Psychiatry. 23(5):1233-1243.
2. Falls DL. (2003) Exp Cell Res. 284(1):14-30.
3. Zhao WJ. et al. (2021) Folia Neuropathol. 59(2):189-197.
4. Czarnek M. et al. (2020) Mol Neurobiol. 57(4):1799-1813.

Synonyms

neuregulin 2, neuregulin2, Neuregulin-2, NTAK, Pro NRG2

Selected Validation Data



Purity of Recombinant Human NRG2 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.

For technical support and original validation data for this product please contact

T: 027-87531629

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.