For Research Use Only

FITC Plus Anti-Human PD-L1/CD274 (29E.2A3) Catalog Number: FITC-65081





Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** FITC-65081 BC069381 Affinity purification GenelD (NCBI): CloneNo.: Size: 100tests, 5 $\,\mu$ l/test 29126 29E.2A3 UNIPROT ID: Excitation/Emission maxima Source: Mouse Q9NZQ7 wavelengths: 495 nm / 524 nm Full Name: Isotype: IgG2b, kappa CD274 molecule **Applications Tested Applications:** FC **Species Specificity:** human **Background Information** Programmed cell death ligand 1 (PD-L1, CD274, or B7-H1), is the first member of B7 family to be discovered. B7 family molecules are type I transmembrane proteins belonging to the immunoglobulin superfamily. In concert with their CD28 family receptors, the B7s are key regulators of the adaptive immune response. PD-L1 is suggested as a negative regulator of T and B cell, and plays important role in mediating tolerance of lymphocytes to self-antigens. It is also involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PD-1-independent manner. Storage Storage: Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 0.09% sodium azide and 0.5% BSA.

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^6 PHA treated (3 days) human PBMCs were surface stained with CL750 Anti-Human CD3 and 5 ul FITC Plus Anti-Human PD-L1 (B7-H1) (FITC-65081, Clone:29E.2A3) (right) or FITC Plus Mouse IgG2b Isotype Control (left). 1x10^6 untreated human PBMCs were surface stained with 5 ul FITC Plus Anti-Human PD-L1 (B7-H1) (FITC-65081, Clone:29E.2A3) (center). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Lymphocytes were gated.

1x10^6 PHA treated (3 days) human PBMCs were surface stained with 5 ul FITC Plus Anti-Human PD-L1 (B7-H1) (FITC-65081, Clone:29E.2A3) (red) or FITC Plus Mouse IgG2b Isotype Control (blue). 1x10^6 untreated human PBMCs were surface stained with 5 ul FITC Plus Anti-Human PD-L1 (B7-H1) (FITC-65081, Clone:29E.2A3) (black dashed). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Lymphocytes were gated.