For Research Use Only

FITC Plus Anti-Human CD28 (CD28.2)

Catalog Number: FITC-65099



Basic Information

Catalog Number:

FITC-65099

Size:

100tests, 5 μ l/test

Source: Mouse

Isotype: IgG1, kappa UNIPROT ID: P10747 Full Name:

CD28 molecule Calculated MW: 220 aa, 25 kDa

BC093698

940

GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000178562

GenBank Accession Number:

Purification Method:

Affinity purification

CloneNo.: CD28.2

Excitation/Emission maxima

wavelengths: 495 nm / 524 nm

Applications

Tested Applications:

FC

Species Specificity:

Human

Background Information

CD28 (T-cell-specific surface glycoprotein CD28), also known as T44 and Tp44, is a 44 kD disulfide-linked homodimeric type I glycoprotein (PMID: 2162180). It is a member of the immunoglobulin superfamily and is expressed on most T lineage cells, NK cell subsets, and plasma cells (PMID: 2162180, 8386518). CD28 may affect in vivo immune responses by functioning both as a cell adhesion molecule linking B and T lymphocytes and as the surface component of a novel signal transduction pathway (PMID: 2162180, 3021470). CD28 binds both CD80 and CD86 with a highly conserved motif MYPPY in the CDR3-like loop (PMID: 15696168, 7964482). CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death (PMID: 1348520).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1X10^6 human PBMCs were surface co-stained with PE Anti-Human CD3 and 5 ul FITC Plus Anti-Human CD28 (FITC-65099, Clone: CD28.2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD28 (FITC-65099, Clone: CD28.2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.