For Research Use Only

FITC Plus Anti-Mouse CD34 Rabbit Recombinant Antibody

Catalog Number: FITC-98069



Basic Information

Catalog Number: GenBank Accession Number: FITC-98069 NM_133654

 Concentration:
 GeneID (NCBI):

 100ug, 500 ug/ml
 12490

 Source:
 UNIPROT ID:

 Rabbit
 Q64314

Isotype: Full Name:
IgG CD34 antigen
Immunogen Catalog Number: Calculated MW:
EG0854 41 kDa

Purification Method: Protein A purification

CloneNo.: 240659B11

Excitation/Emission maxima wavelengths:

495 nm / 524 nm

Applications

Tested Applications:

FC

Species Specificity:

mouse

Background Information

CD34 is a 105- to 120-kDa glycophosphoprotein expressed on the majority of hematopoietic stem/progenitor cells, bone marrow stromal cells, capillary endothelial cells, embryonic fibroblasts, and some nerve tissue. CD34 is a commonly used marker for identifying human hematopoietic stem/progenitor cells and mediates cell adhesion and lymphocyte homing by binding L-selectin and E-selectin ligands. CD34 is also one of the best negative selection markers for characterizing and/or isolating human MSCs from bone marrow and other sources. Along with other positive selection markers (such as CD29, CD44, CD90, CD105 and CD166), negative selection markers (such as CD34 and CD45) are used for MSC identification.


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer.

PBS with 0.09% sodium azide, pH7.3

Selected Validation Data



1x10^6 mouse bone marrow cells were surface stained with 0.25 ug FITC Plus Anti-Mouse CD34 Rabbit RecAb (FITC-98069, Clone: 240659B11), or FITC Plus Rabbit 1gG Isotype Control RecAb (FITC-98136, Clone: 240953C9). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 mouse bone marrow cells were surface stained with 0.25 ug FITC Plus Anti-Mouse CD34 Rabbit RecAb (FITC-98069, Clone: 240659B11) (red), or FITC Plus Rabbit 1gG Isotype Control RecAb (FITC-98136, Clone: 240953C9) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.