For Research Use Only

## FITC Plus Anti-Human Granzyme B Rabbit Recombinant Antibody

Catalog Number: FITC-98070



**Purification Method:** 

Protein A purification

**Basic Information** 

Catalog Number: GenBank Accession Number: FITC-98070 BC030195

Concentration:GeneID (NCBI):CloneNo.:100tests, 5 ul/test3002240761G4

Source: UNIPROT ID: Excitation/Emission maxima
Rabbit P10144 wavelengths:
Isotype: Full Name: 495 nm / 524 nm

IgG granzyme B (granzyme 2, cytotoxic T-

lymphocyte-associated serine

esterase 1)

Calculated MW:
247 aa, 28 kDa

**Applications** 

Tested Applications:

FC (Intra)

Species Specificity:

human

## **Background Information**

GZMB (Granzyme B) is also named as CGL1, CSPB, CTLA1, GRB and belongs to the Granzyme subfamily. This enzyme is necessary for target cell lysis in cell-mediated immune responses. The cytotoxic lymphocyte protease granzyme B (GzmB) can promote apoptosis through direct processing and activation of members of the caspase family. GzmB can also cleave the BH3-only protein, BID, to promote caspase-independent mitochondrial permeabilization (PMID:17283187). GzmB induces lamin B degradation in isolated nuclei less efficiently than GzmA (PMID:11331782). This full-length protein has 2 glycosylation sites and a signal peptide. Unglycosylated human granzyme B is 26 kDa and high mannose glycosylated is 32 kDa and only 32 kDa or smaller forms of granzyme B are accumulated within nuclei (PMID:8626751). GzmB also forms dimers.

Storage


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA

## **Selected Validation Data**



1x10^6 human PBMCs were intracellularly stained with APC Anti-Human CD8a, and 5 ul FITC Plus Anti-Human Granzyme B Rabbit RecAb (FITC-98070, Clone: 240761G4) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1x10^6 human PBMCs were intracellularly stained with 5 ul FITC Plus Anti-Human Granzyme B Rabbit RecAb (FITC-98070, Clone: 240761G4) (red) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).