For Research Use Only

FITC Plus Anti-Mouse PD-1/CD279 Rabbit Recombinant Antibody

Catalog Number: FITC-98080



Basic Information

Catalog Number: GenBank Accession Number: FITC-98080 NM_008798.2

GeneID (NCBI): Concentration: 100ug, 500 µg/ml 18566 **UNIPROT ID:** Source: Rabbit Q02242

Full Name: Isotype:

programmed cell death 1

Calculated MW: Immunogen Catalog Number:

EG0918 32 kDa

Positive Controls:

FC: anti-CD3/CD28 treated mouse splenocytes,

Purification Method:

CloneNo.:

240389B8

suspension

wavelengths:

495 nm / 524 nm

Protein A purification

Recommended Dilutions:

Excitation/Emission maxima

FC: 0.25 ug per 10^6 cells in a 100 μ l

Applications

Tested Applications:

Species Specificity:

mouse

Background Information

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Selected Validation Data



1x10^6 untreated or anti-CD3/CD28 treated mouse splenocytes were surface stained with APC Anti-Mouse CD3 and 0.25 ug FITC Plus Anti-Mouse PD-1/CD279 Rabbit RecAb (FITC-98080, Clone: 24038988). Cells were not fixed.



1x10^6 anti-CD3/CD28 treated mouse splenocytes were surface stained with 0.25 ug FITC Plus Anti-Mouse PD-1/CD279 Rabbit RecAb (FITC-98080, Clone: 240389B8) (red) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9) (blue). 1x10^6 untreated mouse splenocytes were surface stained with 0.25 ug FITC Plus Anti-Mouse PD-1/CD279 Rabbit RecAb (FITC-98080, Clone: 240389B8) (black, dashed). Cells were not fixed.