

For Research Use Only

FITC Plus Anti-Human CD47 Rabbit Recombinant Antibody

Catalog Number: FITC-98131



Basic Information

Catalog Number:

FITC-98131

Concentration:

100 tests, 5 µl/test

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG31499

GenBank Accession Number:

BC010016

GeneID (NCBI):

961

UNIPROT ID:

Q08722

Full Name:

CD47 molecule

Calculated MW:

323 aa, 35 kDa

Purification Method:

Protein A purification

CloneNo.:

240997D8

Excitation/Emission maxima wavelengths:

495 nm / 524 nm

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

CD47, also known as integrin-associated protein (IAP), is a member of the immunoglobulin superfamily containing a five-pass transmembrane attachment. CD47 is heavily glycosylated and widely expressed by hematopoietic and nonhematopoietic cells. CD47 interacts with several membrane integrins and also acts as a receptor for thrombospondin (THBS1). It is involved in a range of cellular processes, including apoptosis, proliferation, adhesion, and migration. CD47 also functions as a ligand for signal regulatory protein- α (SIRP α). Upon binding CD47, SIRP α initiates a signaling cascade that results in the inhibition of phagocytosis.

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ FC receptor blocked human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD47 Rabbit RecAb (FITC-98131, Clone:240997D8) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10⁶ FC receptor blocked human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD47 Rabbit RecAb (FITC-98131, Clone: 240997D8) (red) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.