For Research Use Only

FITC Plus Anti-Human LAIR1 Rabbit Recombinant Antibody

Catalog Number: FITC-98326



Basic Information

Catalog Number: GenBank Accession Number: FITC-98326 BC027899

Concentration: GeneID (NCBI): 100tests, 5 ul/test 3903 **UNIPROT ID:** Source:

Rabbit Q6GTX8 Isotype: Full Name:

IgG leukocyte-associated immunoglobulin-like receptor 1 Immunogen Catalog Number:

EG2329 Calculated MW:

287 aa, 31 kDa

Purification Method:

Protein A purification

CloneNo.: 241805G2

Recommended Dilutions:

FC: 5 ul per 10^6 cells in a 100 µl

suspension

Excitation/Emission maxima

wavelengths: 495 nm / 524 nm

Applications

Tested Applications:

Species Specificity:

human

Positive Controls:

FC: human PBMCs,

Background Information

Leukocyte-associated immunoglobulin-like receptor 1 (LAIR1) is a transmembrane glycoprotein with a single immunoglobulin-like domain and a cytoplasmic tail containing two immune receptor tyrosine-based inhibitory motifs (PMID: 9285412). LAIR1 is expressed on the majority of human PBMCs, including NK, T, B, monocytes, and dendritic cells, as well as the majority of thymocytes (PMID: 10229813). It functions as an inhibitory receptor that plays a constitutive negative regulatory role on cytolytic function of NK cells, B cells and T cells. LAIR1 is a collagen receptor (PMID: 16754721). Tumor-expressed collagens can modulate immune cell function through the inhibitory collagen receptor LAIR1 (PMID: 21955987).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1x10^6 FC receptor blocked human PBMCs were surface stained with CoraLite® Plus 647 Anti-Human CD19, and 5 u IFTC Plus Anti-Human LAIR1 Rabbit RecAb (FITC-98326, Clone: 241805G2) or FITC Plus Rabbit 1gG Isotype Control RecAb (FITC-98136, Clone: 240953C9). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 FC receptor blocked human PBMCs were surface stained with 5 ul FITC Plus Anti-Human LAIR1 Rabbit RecAb (FITC-98326, Clone: 241805G2) (red) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9)(blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.