## For Research Use Only

## MultiPro<sup>TM</sup> 5CFLX Anti-Human Lselectin/CD62L (Polyclonal)




Catalog Number: G26477-1-5C


| Basic Information             | Catalog Number:<br>G26477-1-5C  | GenBank Accession Number:<br>BC020758  | Conjugate:<br>5CFLX  |   |
|-------------------------------|---|--|--|---|
|                               | Size:<br>500ug/mL<br>Source:  | GeneID (NCBI):<br>6402<br>ENSEMBL Gene ID:<br>ENSG00000188404<br>UNIPROT ID:<br>P14151<br>Full Name:<br>MultiPro <sup>TM</sup> 5CFLX Anti-Human L-<br>selectin/CD62L (Polyclonal)  | Full Oligo Sequence:<br>CGGAGATGTGTATAAGAGACAGAGCT<br>TCGCTGAACCGCCCATATAAGAAA<br>Barcode Sequence:<br>AGCTTCGCTGAACCG |   |
|                               |   |  |  | Rabbit<br>Isotype:                          |
|                               | lgG<br>Immunogen Catalog Number:<br>AG24063   |  |  |   |
|                               | Applications  |  |  | Tested Applications:<br>Single Cell (Intra) |
| Species Specificity:<br>Human |   |  |  |   |
| Background Informatio         | CD62E (E-selectin) and CD62P (P-<br>protein of 95-105 kDa on neutropl<br>expressed on the surface of most<br>hematopoietic progenitor cells, a<br>lymphocytes to high endothelial<br>expressed ligand, and is also invo | CD62L, also known as L-selectin or SELL, is a member of the selectin family of adhesion molecules that also include CD62E (E-selectin) and CD62P (P-selectin) (PMID: 2663882, 2473156, 1382078). CD62L is a highly glycosylated protein of 95-105 kDa on neutrophils and 74 kDa on lymphocytes (PMID: 1382078; 1694883, 1695155). CD62L is expressed on the surface of most leukocytes, including lymphocytes, neutrophils, monocytes, eosinophils, hematopoietic progenitor cells, and immature thymocytes (PMID: 1694883, 1688580). It mediates the binding of lymphocytes to high endothelial venules (HEV) of peripheral lymph nodes through interactions with a constitutively expressed ligand, and is also involved in lymphocyte, neutrophil, and monocyte attachment to endothelium at sites of inflammation (PMID: 1382078). |  |   |
| Storage                       | Storage:<br>2-8°C<br>Storage Buffer:<br>PBS with 1mM EDTA and 0.09% so  | odium azide  |  |   |

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





G26477-1-5C was used to stain Resting PBMC and analyzed with 10x Genomics Gene Expression Flex with Feature Barcodes and Multiplexing kit with Fix-Stain protocol. G26477-1-5C was used to stain PBMC under 4hr LPS + TI treatment and analyzed with 10x Genomics Gene Expression Flex with Feature Barcodes and Multiplexing kit with Fix-Stain protocol.