

For Research Use Only

MultiPro™ 5CFLX Anti-Human CD28 (CD28.2)



Catalog Number: G65099-1-5C

Basic Information

Catalog Number: G65099-1-5C	GenBank Accession Number: BC093698	CloneNo.: CD28.2
Size: 500ug/mL	GeneID (NCBI): 940	Conjugate: 5CFLX
Source: Mouse	ENSEMBL Gene ID: ENSG00000178562	Full Oligo Sequence: CGGAGATGTGTATAAGACAGGTTA TCGCAACGTTACCCATATAAGAAA
Isotype: IgG1, kappa	UNIPROT ID: P10747	Barcode Sequence: GTTATCGCAACGTTA
	Full Name: MultiPro™ 5CFLX Anti-Human CD28 (CD28.2)	

Applications

Tested Applications:
Single Cell (Intra), Single Cell

Species Specificity:
Human

Background Information

CD28 (T-cell-specific surface glycoprotein CD28), also known as T44 and Tp44, is a 44 kD disulfide-linked homodimeric type I glycoprotein (PMID: 2162180). It is a member of the immunoglobulin superfamily and is expressed on most T lineage cells, NK cell subsets, and plasma cells (PMID: 2162180, 8386518). CD28 may affect in vivo immune responses by functioning both as a cell adhesion molecule linking B and T lymphocytes and as the surface component of a novel signal transduction pathway (PMID: 2162180, 3021470). CD28 binds both CD80 and CD86 with a highly conserved motif MYPPY in the CDR3-like loop (PMID: 15696168, 7964482). CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death (PMID: 1348520).

Storage

Storage:
2-8°C

Storage Buffer:
PBS with 1mM EDTA and 0.09% sodium azide

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




G65099-1-5C was used to stain Resting PBMC and analyzed with 10x Genomics Gene Expression Flex with Feature Barcodes and Multiplexing kit with Fix-Stain protocol.



G65099-1-5C was used to stain PBMC under 4hr LPS + TI treatment and analyzed with 10x Genomics Gene Expression Flex with Feature Barcodes and Multiplexing kit with Fix-Stain protocol.



G65099-1-5C was used to stain Resting PBMC and analyzed with 10x Genomics Gene Expression Flex with Feature Barcodes and Multiplexing kit with Stain-Fix protocol.



G65099-1-5C was used to stain PBMC under 4hr LPS + TI treatment and analyzed with 10x Genomics Gene Expression Flex with Feature Barcodes and Multiplexing kit with Stain-Fix protocol.