For Research Use Only

MultiProTM 5CFLX Anti-Human CD8 (SK1)



Catalog Number: G65146-1-5C

Basic Information

Catalog Number: GenBank Accession Number: CloneNo.: G65146-1-5C BC025715 SK1 GeneID (NCBI): Conjugate: Size: 500ug/mL 5CFLX

ENSEMBL Gene ID: Full Oligo Sequence: Source:

CGGAGATGTGTATAAGAGACAGATTA Mouse ENSG00000153563 GATTGGTTAACCCCATATAAGAAA Full Name: Isotype:

IgG1, kappa MultiProTM 5CFLX Anti-Human CD8 Barcode Sequence: ATTAGATTGGTTAAC

(SK1)

Applications

Tested Applications: Single Cell (Intra) Species Specificity: Human

Background Information


 ${\sf CD8}\ is\ a\ transmembrane\ glycoprotein\ composed\ of\ two\ disulfide-linked\ chains.\ It\ can\ be\ present\ as\ a\ homodimer\ of\ composed\ of\ two\ disulfide-linked\ chains.\ It\ can\ be\ present\ as\ a\ homodimer\ of\ composed\ of\ two\ disulfide-linked\ chains.\ It\ can\ be\ present\ as\ a\ homodimer\ of\ composed\ of\ two\ disulfide-linked\ chains.\ It\ can\ be\ present\ as\ a\ homodimer\ of\ composed\ of\ two\ disulfide-linked\ chains.\ It\ can\ be\ present\ as\ a\ homodimer\ of\ composed\ of\ two\ disulfide-linked\ chains.\ It\ can\ be\ present\ as\ a\ homodimer\ of\ composed\ of\ two\ disulfide-linked\ chains.\ It\ can\ be\ present\ as\ a\ homodimer\ of\ composed\ o$ CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α α homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) γ δ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Storage


Storage: 2-8°C Storage Buffer:

PBS with 1mM EDTA and 0.09% sodium azide

Selected Validation Data



G65146-1-5C was used to stain Resting PBMC and analyzed with 10x Genomics Gene Expression Flex with Feature Barcodes and Multiplexing kit with Fix-Stain protocol.



G65146-1-5C was used to stain PBMC under 4hr LPS + TI treatment and analyzed with 10x Genomics Gene Expression Flex with Feature Barcodes and Multiplexing kit with Fix-Stain protocol.