For Research Use Only

HRP-conjugated PCNA Monoclonal antibody

www.ptglab.com

Purification Method:

CloneNo.:

10D10E11

Protein G purification

Recommended Dilutions:

WB 1:5000-1:50000

Catalog Number: HRP-60097

Featured Product

2 Publications

Basic Information

Applications

Catalog Number: HRP-60097 Size:

1000 µg/ml Source: Mouse Isotype:

lgG1 Immunogen Catalog Number:

AG7416

Tested Applications:

WB

Cited Applications:

WB

Species Specificity: human, mouse, pig, rat Cited Species: human, mouse

GenBank Accession Number:

BC000491 GeneID (NCBI):

UNIPROT ID: P12004 Full Name:

proliferating cell nuclear antigen

Calculated MW: 29 kDa/31 kDa Observed MW:

36-38 kDa

Positive Controls:

WB: HEK-293 cells,

Background Information

Proliferating Cell Nuclear Antigen, commonly known as PCNA, is a protein that acts as a processivity factor for DNA polymerase δ in eukaryotic cells. This protein is an auxiliary protein of DNA polymerase delta and is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand. PCNA induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic-apyrimidinic (AP) endonuclease, APEX2 activities. It has to be loaded onto DNA in order to be able to stimulate APEX2. PCNA protein is highly conserved during evolution; the deduced amino acid sequences of rat and human differ by only 4 of 261 amino acids. PCNA has been used as loading control for proliferating cells. The calculated molecular weight of PCNA is 29 kDa, but modified PCNA is 36kDa (PMID: 1358458).

Notable Publications

Author	Pubmed ID	Journal	Application
Ha T Nguyen	36468795	Mol Reprod Dev	WB
Shengjie Zhu	39638770	Phytother Res	WB


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with HRP-60097 (PCNA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.