For Research Use Only


HRP-conjugated 6*His, His-Tag Monoclonal antibody Catalog Number:HRP-66005 166 Publications



Basic Information	Catalog Number: HRP-66005	GenBank Acc GeneID (NCB	ession Number:	Purification Method: Protein G purification	
	Concentration:	Full Name:		CloneNo.:	
	1000 µg/ml Source: Mouse	Calculated MW: 0.84 kDa	1W:	187G5 Recommended Dilutions: WB 1:5000-1:50000	
	lsotype: lgG1				
Applications	Tested Applications:		Positive Controls:		
	WB WB : Recombinant protein protein,				
	Cited Applications: WB, IF, IP, CoIP, ELISA, Blocking assay				
	Species Specificity: recombinant protein				
Background Information	Protein tags are protein or peptide sequences located either on the C- or N- terminal of the target protein, which facilitates one or several of the following characteristics: solubility, detection, purification, localization and expression. His-tag is often used for affinity purification and binding assays. Expressed His-tagged proteins can be purified and detected easily because the string of histidine residues binds to several types of immobilized metal ions, including nickel, cobalt and copper, under specific buffer conditions. This His-tag antibody is a useful tool for monitoring of the His-tagged proteins, and recognizes His-tags placed at N-terminal, C-terminal, and internal regions of fusion proteins expressed in bacteria, insect, and mammalian cells. This antibody is conjugated with His-				
	Note: Do not add Azium (Sodium Azide or Smite) into the dilution buffer. Azium is the HRP inhibitor which decrease the enzymatic activity of HRP.				
	A. (1	Pubmed ID	Journal	Application	
Notable Publications	Author				
Notable Publications	Author Antti Kulmala	36178799	ACS Synth Biol	WB	
Notable Publications		36178799 33003404	ACS Synth Biol Cells	WB	
Notable Publications	Antti Kulmala		•	WB	

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Recombinant protein were subjected to SDS PAGE followed by western blot with HRP-66005 (6°His, His-Tag antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.