

**HumanKine<sup>®</sup> PDGF $\alpha\alpha$  (Recombinant Human)**



Animal Component-Free	Human cell expressed	Tag-Free	Endotoxin Free
-----------------------	----------------------	----------	----------------

**Product Description**


Animal-free Recombinant Human PDGF $\alpha\alpha$  is expressed in human 293 cells as a mixture of pro-dimers, pro-mature and mature dimers with an apparent molecular mass of 35 to 45 kDa. The cytokine is produced in a human cell expression system with serum free, chemically defined media.

Alternative Names	PDGF 1, PDGF A, PDGF subunit A, PDGF1, PDGFA, PDGF $\alpha\alpha$
Source	Human Embryonic Kidney cells (HEK293). HEK293-derived PDGF $\alpha\alpha$ protein

**Specifications**


Test	Method	Specification
Activity	Dose-dependent stimulation of the proliferation of 3T3 cells. Dose-dependent stimulation of the proliferation of NIH 3T3 cells with low serum medium containing BSA and Dexamethasone	Typically $\leq 10$ ng/mL in 3T3 cell, 150 - 900 ng/mL in NIH 3T3
Molecular Mass	SDS-PAGE	19 and 26 reduced, 35, 40 and 45 kDa non-reduced, homodimer, glycosylated
Purity	SDS-PAGE	>95%
Endotoxin	LAL	<1 EU/ $\mu$ g

**Activity Data**



Recombinant human PDGF $\alpha\alpha$  (HZ-1215) stimulates dose-dependent proliferation of the NIH/3T3 mouse fibroblast cell line. Viable cell number was quantitatively assessed by Prestoblu<sup>e</sup> Cell Viability Reagent. NIH/3T3 cells were serum starved with 0.1% FBS for 24 hours before treatment with increasing concentrations of recombinant human PDGF $\alpha\alpha$  for 48hrs. The EC<sub>50</sub> was

**SDS-PAGE**



Preparation	
Shipping Temperature	ambient temperature
Formulation	10 mM HOAc, See Certificate of Analysis for details
Reconstitution	Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein to 0.2 mg/mL in sterile 10 mM HOAc containing 0.1% endotoxin-free recombinant human serum albumin (HSA). Gently swirl or tap vial to mix.

Stability and Storage	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
	Lyophilized	Room Temperature	2 weeks
	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
Avoid repeated freeze-thaw cycles.			

[www.ptglab.com](http://www.ptglab.com)

Document #: FR-QA118-101 Rev 0  
Data Sheet Version #: 1

**Proteintech Group, Inc.**  
5500 Pearl Street, Suite 400 Rosemont, IL 60612  
t: 1-888-478-4522; f: 1-312-455-8408  
Email: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)