

Catalog Number: HZ-1321

Data Sheet



HumanKine® IL-12 beta (Recombinant Human)

Animal Component-Free

Human cell expressed

Tag-Free


Endotoxin Free

Product Description

IL-12p40, also known as IL-12B, is a subunit of both the IL-12 and IL-23 cytokines. It is not thought to be bioactive on its own, but it has been shown to regulate IL-12 activity in antagonistic manner by competitively binding with the IL-12 receptor. IL-12p40 also functions as a chemoattractant molecule for macrophage infiltration into tumor sites. There has additionally been evidence to suggest that IL-12p40 mediates a protective immune response against mycobacterial infection by serving as a necessary component for granuloma formation and promoting Th1 effector cell functions (PMID: 17126601, 10453037, 11739515).

Alternative Names	Interleukin-12 subunit beta (IL-12B) Cytotoxic lymphocyte maturation factor (CLMF p40), IL-12 subunit p40, NK cell stimulatory factor chain 2 (NKSF2)
Source	Human Embryonic Kidney cells (HEK293). HEK293-derived IL-12 beta protein

Specifications					
Test Method Specification					
Activity	Inquire	Inquire			
Molecular Mass	SDS-PAGE	37-50 kDa reduced, 37 to 47, 65 to 95, and 125 kDa non-reduced, monomer, glycosylated			
Purity	SDS-PAGE	> 95%			
Endotoxin	LAL	< 0.1 EU/ug			



www.ptglab.com

Document #: FR-QA118-101 Rev 0
Data Sheet Version #: 1

Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com

Preparation				
Shipping Temperature	ambient temperature			
Formulation	1x PBS pH 7.4			
Reconstitution	Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein to 0.2 mg/mL in sterile 1x PBS pH 7.4 containing 0.1% endotoxin-free recombinant human serum albumin (HSA). Gently swirl or tap vial to mix.			

	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
Stability and Storage	Lyophilized	Room Temperature	2 weeks
_	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
		Avoid repeated freeze-thaw cycles.	

www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com