## For Research Use Only FlexAble CoraLite® Plus 750 Antibody Labeling Kit for Mouse IgG2a Catalog Number: KFA044



www.ptgcn.com

## **Product Information**

| Product name                             | FlexAble CoraLite® Plus 750 Antibody Labeling Kit for Mouse IgG2a |
|--|---|
| Assay type                               | Antibody labeling   |
| Tested applications                      | IF, FC, WB  |
| Species Reactivity                       | Mouse IgG2a   |
| Antibody amount per labeling reaction    | 0.5 µg antibody   |
| Conjugate                                | CoraLite® Plus 750  |
| Excitation / Emission maxima wavelengths | 755 nm / 780 nm   |

10 rxns

50 rxns

4×50 rxns

## **Kit Components**

|                   | CoraLite® Plus 750 FlexLinker for Mouse IgG2a   | 10 µ L  | 50 µ L          | 4×50 µL         |  |  |
|-------------------|---|---------|-----------------|-----------------|--|--|
|                   | FlexQuencher for Mouse IgG2a  | 20 µ L  | 100 µL          | <b>4×100</b> μL |  |  |
|                   | FlexBuffer  | 100 µ L | 500 μL          | <b>4×500</b> μL |  |  |
| 包装规格              | 10/50/4x 50 reactions   |         |                 |                 |  |  |
| Storage Condition | Store for 1 year at -20°C or for 6 months at +4°C upon receipt. Avoid exposure to light.  |         |                 |                 |  |  |
| FAQ               | Q: What are the FlexLinker, FlexQuencher and FlexBuffer?<br>A: The FlexLinker is a small polypeptide to which dyes are covalently conjugated that can label unconjugated<br>primary antibodies. The FlexQuencher is an Fc-containing fragment that neutralizes the excess FlexLinker.<br>The FlexBuffer is a PBS-based buffer.  |         |                 |                 |  |  |
|                   | Q: What is the largest quantity I can label?<br>A: With a standard kit size (50 reactions), you can label 25 µg of one antibody or up to 50 different antibodies.<br>You can easily scale up the antibody amount per labeling approach.   |         |                 |                 |  |  |
|                   | Q: What is the lowest concentration of my primary antibody that I can use?<br>A: Our protocol uses 0.5 μg of primary antibody in 7 μL, which ends up at 0.07 mg/mL. If the concentration of<br>your antibody is lower, you can also use a larger volume than 7 μL.  |         |                 |                 |  |  |
|                   | Q: Can I label primary antibodies stored in BSA, glycerol, Tris buffer and/or preservatives?<br>A: Yes, FlexAble Antibody Labeling Kits have been validated with carriers and amine buffers. Neither BSA nor<br>amine buffers, in any chosen concentration, interfere with the labeling. 50% glycerol as well as preservatives<br>like sodium azide are also compatible with the kit.   |         |                 |                 |  |  |
|                   | Q: How many different primary antibodies can I label with one I<br>A: You can label up to 50 different antibodies with our FlexAble<br>FlexAble 10 rxn Kit.   |         | up to 10 antibo | odies with our  |  |  |
|                   | Q: Will I observe cross-reactivity/leaking when I use two FlexAble-labeled antibodies from the same species<br>during multiplexing?<br>A: FlexAble labels primary antibodies with a high affinity FlexLinker. Dissociation of FlexLinker from one<br>antibody and association to another antibody is rare. If you observe leaking, we recommend adding more<br>FlexQuencher to remove unbound FlexLinker, or you can try sequential staining of the labeled antibodies. |         |                 |                 |  |  |
|                   | More FAQs   |         |                 |                 |  |  |

Component


## Validation Data




PFA-fixed HeLa cells were stained with anti-GNL3 (67169-1-1g) labeled with FlexAble Coralite® Plus 405 Kit (KFA046, cyan) and anti-Tubulin (66240-1-1g) labeled with FlexAble Coralite® Plus 750 Kit (KFA044, grey). Epifluorescence images were acquired with a 20x objective and post-processed.



PFA-fixed HeLa cells were stained with anti-Tubulin (66240-1-1g) labeled with FlexAble CoraLite® Plus 750 Kit (KFA044, grey) and Sytox Deep Red (cell nuclei, magenta). Epifluorescence images were acquired with a 20x objective and post-processed.



Flow cytometry of PBMC. 1X10^6 human peripheral blood mononuclear cells (PBMCs) were stained with anti-human CD45 (clone HI 30, 65109-1-1g) labeled with FlexAble CoraLite® Plus 405 Kit (KFA026) and anti-human CD3 (clone OKT3, 65133-1-1g) labeled with FlexAble CoraLite® Plus 750 Kit (KFA044).



Flow cytometry of PBMC. 1X10<sup>^6</sup> human peripheral blood mononuclear cells (PBMCs) were stained with anti-human CD3 (clone UCHT1, 65151-1-1g) labeled with FlexAble Coralite® Plus 405 Kit (KFA026) and anti-human CD8 (clone UCHT4, 65204-1-1g) labeled with FlexAble Coralite® Plus 750 Kit (KFA044).