



IHCeasy EXOSC10 Ready-To-Use IHC Kit

Catalog Number: KHC1665

General Information

Sample type: FFPE tissue Cited sample type: Reactivity: Human, Mouse, Rat Cited Reactivity: Assay type: Immunohistochemistry Primary antibody type: Rabbit Polyclonal

Secondary antibody type: Polymer-HRP-Goat anti-Rabbit

Kit Component

Antigen Retrieval Buffer 100 mL	50×
Washing Buffer 100 mL × 2	20×
Blocking Buffer 5 mL	RTU
Primary Antibody 5 mL	RTU
Secondary Antibody 5 mL	RTU
Chromogen Component A 0.2 mL	RTU
Chromogen Component B 4 mL	RTU
Signal Enhancer 5 mL	RTU
Counter Staining Reagent 5 mL	RTU
Mounting Media 5 mL	RTU
Control Slide 1 slide (Optional)	FFPE
Datasheet 1 Copy	
Manual 1 Copy	

Storage Instructions

All the reagents are stored at 2-8°C. The kit is stable for 6 months from the date of receipt.


Background

About 50% of patients with polymyositis/scleroderma (PM-Scl) overlap syndrome are reported to have autoantibodies to a neuclear/nucleolar particle termed PM-Scl. Exosome component 10 (EXOSC10), also named autoantigen PM/Scl 2, is the 100 kDa antigen component of PM-Scl and is recognized by most sera of PM-Scl paitents. EXOSC10 is strongly enriched in the nucleolus and a small amount has been found in cytoplasm supporting the existence of a nucleolar RNA exosome complex form. As a putative catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease activity, EXOSC10 participates in a multitude of cellular RNA processing and degradation events.


Synonyms

Autoantigen PM/Scl 2, EXOSC10, exosome component 10, p2, p3, p4, PM Scl, PM/Scl 100, PMSCL, PMSCL2, RRP6, Rrp6p


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using KHC1665 (EXOSC10 IHC Kit).



Immunohistochemical analysis of paraffinembedded mouse skin tissue slide using KHC1665 (EXOSC10 IHC Kit).



Immunohistochemical analysis of paraffinembedded rat skin tissue slide using KHC1665 (EXOSC10 IHC Kit).