



IHCeasy SUPT16H Ready-To-Use IHC Kit

Catalog Number: KHC2026

General Information

Sample type: FFPE tissue Cited sample type: Reactivity: Human, Mouse, Rat Cited Reactivity: Assay type: Immunohistochemistry Primary antibody type: Rabbit Polyclonal

Secondary antibody type: Polymer-HRP-Goat anti-Rabbit

Kit Component

| Component | Size | Concentration |
|--------------------------|--------------------|---------------|
| Antigen Retrieval Buffer | 100 mL | 50× |
| Washing Buffer | 100 mL ×2 | 20× |
| Blocking Buffer | 5 mL | RTU |
| Primary Antibody | 5 mL | RTU |
| Secondary Antibody | 5 mL | RTU |
| Chromogen Component A | 0.2 mL | RTU |
| Chromogen Component B | 4 mL | RTU |
| Signal Enhancer | 5 mL | RTU |
| Counter Staining Reagent | 5 mL | RTU |
| Mounting Media | 5 mL | RTU |
| Control Slide | 1 slide (Optional) | FFPE |
| Datasheet | 1 Copy | |
| Manual | 1 Copy | |

Storage Instructions

All the reagents are stored at 2-8°C. The kit is stable for 6 months from the date of receipt.


Background

SUPT16H, also named as FACT140, FACTP140, SPT16 and CDC68, belongs to the peptidase M24 family and SPT16 subfamily. SUPT16H is a component of the FACT complex, a general chromatin factor that acts to reorganize nucleosomes. The FACT complex is involved in multiple processes that require DNA as a template such as mRNA elongation, DNA replication and DNA repair. During transcription elongation the FACT complex acts as a histone chaperone that both destabilizes and restores nucleosomal structure. It facilitates the passage of RNA polymerase II and transcription by promoting the dissociation of one histone H2A-H2B dimer from the nucleosome, then subsequently promotes the reestablishment of the nucleosome following the passage of RNA polymerase II. The FACT complex is probably also involved in phosphorylation of 'Ser-392' of p53/TP53 via its association with CK2 (casein kinase II). It also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene. Primary antibody in this kit is specific to SUPT16H.


Synonyms

FACTp 140, FACT 140, FACT 140 kDa subunit, Facilitates chromatin transcription complex subunit SPT16, Chromatin-specific transcription elongation factor 140 kDa subunit


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human placenta tissue slide using KHC2026 (SUPT16H IHC Kit).



Immunohistochemical analysis of paraffinembedded human skin cancer tissue slide using KHC2026 (SUPT16H IHC Kit).




Immunohistochemical analysis of paraffinembedded mouse intestine tissue slide using KHC2026 (SUPT16H IHC Kit).




Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using KHC2026 (SUPT16H IHC Kit).



Immunohistochemical analysis of paraffinembedded rat skin tissue slide using KHC2026 (SUPT16H IHC Kit).



Immunohistochemical analysis of paraffinembedded rat small intestine tissue slide using KHC2026 (SUPT16H IHC Kit).



Immunohistochemical analysis of paraffinembedded rat testis tissue slide using KHC2026 (SUPT16H IHC Kit).