

IHC*easy* PRKCQ Ready-To-Use IHC Kit

Catalog Number: **KHC2392**

General Information

Sample type:
FFPE tissue
Cited sample type:
Reactivity:
Human, Mouse
Cited Reactivity:

Assay type:
Immunohistochemistry
Primary antibody type:
Rabbit Polyclonal
Secondary antibody type:
Polymer-HRP-Goat anti-Rabbit

Kit Component

Component	Size	Concentration
Antigen Retrieval Buffer	100 mL	50×
Washing Buffer	100 mL ×2	20×
Blocking Buffer	5 mL	RTU
Primary Antibody	5 mL	RTU
Secondary Antibody	5 mL	RTU
Chromogen Component A	0.2 mL	RTU
Chromogen Component B	4 mL	RTU
Signal Enhancer	5 mL	RTU
Counter Staining Reagent	5 mL	RTU
Mounting Media	5 mL	RTU
Control Slide	1 slide (Optional)	FFPE
Datasheet	1 Copy	
Manual	1 Copy	

Storage Instructions

All the reagents are stored at 2-8°C. The kit is stable for 6 months from the date of receipt.


Background

PRKCQ, also named as PRKCT and nPKC-theta, belongs to the protein kinase superfamily, AGC Ser/Thr protein kinase family and PKC subfamily. PRKCQ catalyzes the reaction: ATP + a protein = ADP + a phosphoprotein. PRKCQ is a calcium-independent, phospholipid-dependent, serine- and threonine-specific enzyme. It is essential for T-cell receptor (TCR)-mediated T-cell activation, but it is dispensable during TCR-dependent thymocyte development. PRKCQ links the TCR signaling complex to the activation of NF-kappa-B in mature T lymphocytes. It is required for interleukin-2 (IL2) production. PKC is activated by diacylglycerol which in turn phosphorylates a range of cellular proteins. PKC also serves as the receptor for phorbol esters, a class of tumor promoters.


Synonyms

Protein kinase C theta type, protein kinase C, PRKCT, nPKC-theta, EC:2.7.11.13

Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human thyroid cancer tissue slide using KHC2392 (PRKCQ IHC Kit).



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using KHC2392 (PRKCQ IHC Kit).