



# IHCeasy MAP1S Ready-To-Use IHC Kit

Catalog Number: KHC2571

General Information

Sample type: FFPE tissue Cited sample type: Reactivity: Human, Mouse, Rat Cited Reactivity: Assay type: Immunohistochemistry Primary antibody type: Rabbit Polyclonal

Secondary antibody type: Polymer-HRP-Goat anti-Rabbit

#### Kit Component

| Component                | Size               | Concentration |
|--------------------------|--------------------|---------------|
| Antigen Retrieval Buffer | 100 mL             | 50×           |
| Washing Buffer           | 100 mL ×2          | 20×           |
| Blocking Buffer          | 5 mL               | RTU           |
| Primary Antibody         | 5 mL               | RTU           |
| Secondary Antibody       | 5 mL               | RTU           |
| Chromogen Component A    | 0.2 mL             | RTU           |
| Chromogen Component B    | 4 mL               | RTU           |
| Signal Enhancer          | 5 mL               | RTU           |
| Counter Staining Reagent | 5 mL               | RTU           |
| Mounting Media           | 5 mL               | RTU           |
| Control Slide            | 1 slide (Optional) | FFPE          |
| Datasheet                | 1 Copy             |               |
| Manual                   | 1 Copy             |               |

## Storage Instructions

All the reagents are stored at 2-8°C. The kit is stable for 6 months from the date of receipt.


### Background

MAP1S (also known as C19ORF5 or VCY2IP1) is a novel member of the microtubule-associated protein 1 family and a homologue of the exclusively neuronal distributed microtubule-associated protein 1A and 1B (MAP1A/B). In contrast to MAP1A and MAP1B, MAP1S is expressed in a wide range of tissues in addition to neurons. MAP1S is synthesized as a precursor protein that is partially cleaved into heavy and light chains in a tissue-specific manner. In addition, a short chain isoform may be induced under prolonged mitotic arrest or inhibition of the 26S proteasome. Recently it has been reported that the short chain isoform associates with mitochondria in addition to microtubules and causes irreversible aggregation of dysfunctional mitochondria resulting in cell death.


### **Synonyms**

MAP1S,BPY2-interacting protein 1,BPY2IP1,C19orf5,M


#### **Selected Validation Data**




Immunohistochemical analysis of paraffinembedded human placenta tissue slide using KHC2571 (MAP1SIHC Kit).




Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using KHC2571 (MAP1SIHC Kit).




Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using KHC2571 (MAP1SIHC Kit).




Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using KHC2571 (MAP1S IHC Kit).



Immunohistochemical analysis of paraffinembedded mouse lung tissue slide using KHC2571 (MAP1SIHC Kit).



Immunohistochemical analysis of paraffinembedded rat kidney tissue slide using KHC2571 (MAP1SIHC Kit).



Immunohistochemical analysis of paraffinembedded rat lung tissue slide using KHC2571 (MAP1S IHC Kit).