

chromotek® Myc-Trap® 2.0 Agarose, Kit for Immunoprecipitation

Product Code: yt2ak

Product Information

Description: The ChromoTek Myc-Trap® Agarose, Kit for Immunoprecipitation consists of an anti-Myc NANOBODY®/VHH, which is coupled to agarose beads. It also contains lysis, wash, and elution buffers that can be used for the immunoprecipitation of Myc-fusion proteins from cell extracts of various organisms.

Applications: IP, Co-IP

Specificity/Target: Binds specifically to the Myc-tag (sequence EQKLISEEDL) at the N-terminus, C-terminus, or internal site of the fusion protein. Endogenous c-myc is NOT bound.

Binding capacity: 25 µg of recombinant myc-tagged protein (~30 kDa) per 25 µL bead slurry

Bead Size: 90 µm (cross-linked 4 % agarose beads)

Elution Buffer: 2x SDS-sample buffer (Lämmli), 200 mM glycine pH 2.5,
0.1 mg/ml ChromoTek 2x Myc-peptide (2yp) in PBS pH 7.4

Wash Buffer Compatibility: 1M NaCl, 5 mM DTT, 5 mM β-mercaptoethanol, 5 mM TCEP, 2% NP40,
2% Triton X-100, 0.2% SDS, 2 M Urea

Type: Nanobody

Class: Recombinant

Host: Alpaca

Shipment: Shipped at ambient temperature

Storage Buffer: 20 % ethanol


Storage Condition: Upon receipt store at +4°C. Do not freeze!

Stability: Stable for 1 year upon receipt

chromotek® Myc-Trap® 2.0 Agarose, Kit for Immunoprecipitation

Product Code: yt2ak

Selected Validation Data



Immunoprecipitation of GFP-Myc fusion protein from HEK293T cells using Myc-Trap 2.0 Agarose. IP was done using both untransfected (Mock) and transfected (GFP-myc) cells. I: Input, F: Flow-through, B: Bound.

Kit Components

Component	Composition	Quantity
Myc-Trap® 2.0 Agarose	Anti-Myc VHH cross-linked with agarose beads	20 rxns (500 ul slurry)
Lysis Buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.5 % Nonidet™ P40 Substitute, 0.09 % sodium azide	30 mL
RIPA Buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.1 % SDS, 1 % Triton™ X-100, 1 % Deoxycholate, 0.09 % sodium azide	30 mL
Dilution Buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.018 % sodium azide	50 mL (after dilution with 40 mL water)
Wash Buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.05 % Nonidet™ P40 Substitute, 0.5 mM EDTA, 0.018 % sodium azide	50 mL (after dilution with 40 mL water)
Acidic elution buffer	200 mM glycine pH 2.5	3 x 1 mL

Note: Use your equivalent cell lysis buffer for other cell types like yeast, plants, insects, bacteria. Consider using a Wash buffer without detergent for Co-IP.

chromotek® Myc-Trap® 2.0 Agarose, Kit for Immunoprecipitation

Product Code: yt2ak

Required Buffer Solutions

Buffer	Composition
2x SDS-sample buffer	120 mM Tris/Cl pH 6.8, 20% glycerol, 4% SDS, 0.04% bromophenol blue, 10% β-mercaptoethanol
Neutralization buffer	1M Tris pH 10.4 (adjust the pH at +4°C
Peptide elution buffer	0.1 mg/ml 2x Myc-peptide (2yp) reconstituted in PBS

Product Sizes

Product	Product Code	Size
Myc-Trap® 2.0 Agarose, Kit for Immunoprecipitation	yt2ak-20	20 reactions including buffers

chromotek® Myc-Trap® 2.0 Agarose, Kit for Immunoprecipitation

Product Code: yt2ak

Protocol at a glance

General


- Perform all steps at 4°C
- Use your preferred cell lysis buffer and cell lysis conditions

Cell Lysis



- Use 10^6 - 10^7 cells and 200 μ L Lysis buffer.
- Perform cell lysis and clear lysate
- Mix 200 μ L cleared lysate with 300 μ L dilution buffer.

Bead Equilibration




- Transfer 25 μ L bead slurry into a 1.5 mL tube
- Equilibrate beads 3x with 500 μ L dilution buffer

Protein binding



- Add 500 μ L diluted lysate to beads
- Rotate end-over-end for 1 hour at 4°C.

Washing



- Wash beads 3x with 500 μ L wash buffer
- Transfer beads to a new tube during the last washing step

Elution with SDS-sample buffer



- Resuspend beads in 80 μ L 2x SDS-sample buffer
- Boil beads for 5 min at 95°C
- Analyze the supernatant in SDS-PAGE/ Western Blot

chromotek® Myc-Trap® 2.0 Agarose, Kit for Immunoprecipitation

Product Code: yt2ak

Immunoprecipitation Protocol

Cell Material

The following protocol describes the preparation of a mammalian cell lysate.

For other types of cells, we recommend using 500 µg of cell extract and start the protocol with step Bead equilibration.

Mammalian Cell Lysis

Note: Harvesting of cells and cell lysis should be performed with ice-cold buffers. We strongly recommend adding protease inhibitors to the Lysis buffer to prevent degradation of your target protein and its binding partners.

For one immunoprecipitation reaction, we recommend using $\sim 10^6$ - 10^7 cells.

1. Choice of lysis buffer:
 - a. For cytoplasmic proteins, resuspend the cell pellet in 200 µL ice-cold Lysis buffer by pipetting up and down. Supplement Lysis buffer with protease inhibitor cocktail and 1 mM PMSF (not included).
 - b. For nuclear/chromatin proteins, resuspend cell pellet in 200 µL ice-cold RIPA buffer supplemented with DNase I (f.c. 75-150 Kunitz U/mL), MgCl_2 (f.c. 2.5 mM), protease inhibitor cocktail and PMSF (f.c. 1 mM) (not included).
2. Place the tube on ice for 30 min and extensively pipette the suspension every 10 min.
3. Centrifuge cell lysate at 17,000x g for 10 min at +4°C. Transfer cleared lysate (supernatant) to a pre-cooled tube and add 300 µL Dilution buffer supplemented with 1 mM PMSF and protease inhibitor cocktail (not included). If required, save 50 µL of diluted lysate for further analysis (input fraction).

Bead Equilibration

1. Resuspend the beads by gently pipetting them up and down or by inverting the tube. Do not vortex the beads!
2. Transfer 25 µL of bead slurry into a 1.5 mL reaction tube.
3. Add 500 µL ice-cold Dilution buffer.
4. Sediment the beads by centrifugation at 2,500x g for 5 min at +4°C. Discard the supernatant.

Note: Alternatively, ChromoTek Spin columns (sct-10; -20; -50) can be used to equilibrate the beads.

Protein Binding

1. Add diluted lysate to the equilibrated beads.
2. Rotate end-over-end for 1 hour at +4°C.

chromotek® Myc-Trap® 2.0 Agarose, Kit for Immunoprecipitation

Product Code: yt2ak

Washing

1. Sediment the beads by centrifugation at 2,500x g for 5 min at +4°C.
2. If required, save 50 µL of supernatant for further analysis (flow-through/non-bound fraction).
3. Discard remaining supernatant.
4. Resuspend beads in 500 µL Wash buffer.
5. Sediment the beads by centrifugation at 2,500x g for 5 min at +4°C. Discard remaining supernatant.
6. Repeat this step at least twice.
7. During the last washing step, transfer the beads to a new tube.

Optional: To increase stringency of the Wash buffer, test various salt concentrations e.g. 150-500 mM, and/or add a non-ionic detergent e.g. Triton™ X-100 (see Wash buffer compatibility table for maximal concentrations).

Note: Alternatively, ChromoTek Spin columns (sct-10; -20; -50) can be used to wash the beads.

Elution with 2x SDS-sample buffer (Laemmli)

1. Remove the remaining supernatant.
2. Resuspend beads in 80 µL 2x SDS-sample buffer.
3. Boil beads for 5 min at +95°C to dissociate immunocomplexes from beads.
4. Sediment the beads by centrifugation at 2,500x g for 2 min at +4°C.
5. Analyze the supernatant in SDS-PAGE / Western Blot.

Note: For Western blot detection we recommend ChromoTek rat anti Myc-tag Monoclonal antibody (9E1) (Proteintech 9e1) and HRP-conjugated Goat Anti-Rat IgG(H+L) (Proteintech SA00001-15).

Elution with Acidic Elution Buffer

1. Remove the remaining supernatant.
2. Add 50-100 µL Acidic elution buffer and constantly pipette up and down for 30-60 sec at +4°C or room temperature.
3. Sediment the beads by centrifugation at 2,500x g for 2 min at +4°C.
4. Transfer the supernatant to a new tube.
5. Immediately neutralize the eluate fraction with 5-10 µL Neutralization buffer.
6. Repeat this step at least once to increase elution efficiency.

Note: Elution at room temperature is more efficient than elution at +4°C. Prewarm buffers for elution at room temperature.

chromotek® Myc-Trap® 2.0 Agarose, Kit for Immunoprecipitation

Product Code: yt2ak

Elution with 2x Myc-peptide

1. Reconstitute 1 mg ChromoTek 2x Myc-peptide (2yp) in 1000 µl PBS pH 7.4, which results in a final concentration of 1 mg/ml (420 µM). Vortex for 1 min to dissolve the powder. Aliquot upon reconstitution. Dilute 2x Myc-peptide stock to 0.1 mg/ml (42 µM) in PBS for elution.
2. Remove the remaining supernatant from the beads.
3. Add 80 µL 2x Myc-peptide (0.1 mg/ml) and mix using a pipette.
4. Incubate at room temperature for 5-15 min under regular pipetting to ensure thorough mixing.
5. Sediment the beads by centrifugation at 2,500x g for 2 min at +4°C.
6. Transfer the supernatant to a new tube.
7. Repeat this step at least once to increase elution efficiency.

Note: Alternatively, ChromoTek Spin columns (sct-10; -20; -50) can be used to separate the beads.

Related Products

Product	Code
2x Myc-peptide	2yp
Myc-Trap® 2.0 Agarose	yt2a
Myc-Trap® 2.0 Magnetic Agarose	yt2ma
Myc-Trap® 2.0 Magnetic Agarose Kit	yt2mak

Contact

Proteintech North America (HQ)

Proteintech Group, Inc.
5500 Pearl Street, Suite 400
Rosemont, IL 60018 USA

1-888-472-4522

proteintech@ptglab.com

Proteintech Europe

Transmission (6th Fl)
6 Atherton Street
M3 3GS, Manchester, UK

+44 161 839 3007

europe@ptglab.com

chromotek® Myc-Trap® 2.0 Agarose, Kit for Immunoprecipitation

Product Code: yt2ak

Proteintech Germany

Fraunhoferstr. 1
82152, Planegg-Martinsreid
Germany

+49 89 124 148 850
germany@ptglab.com

Disclaimer

For research use only (RUO), not for diagnostic or therapeutic use.

ChromoTek® and Myc-Trap® are registered trademarks of Proteintech Germany GmbH, part of Proteintech Group.
NANOBODY® is a registered trademark of Ablynx N.V.