For Research Use Only

PE Anti-Human CD11b (ICRF44)

Catalog Number:PE-65116 2 Publications



Basic Information

Catalog Number:

PE-65116

Size:

100tests, 5 ul/test

Source: Mouse

Isotype: IgG1, kappa

UNIPROT ID: P11215 Full Name:

integrin, alpha M (complement component 3 receptor 3 subunit)

GenBank Accession Number:

BC096346

3684

GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000169896

Calculated MW: 1152 aa, 127 kDa **Purification Method:**

Affinity purification

CloneNo.: ICRF44

Excitation/Emission maxima

wavelengths:

496 nm, 565 nm / 578 nm

Applications

Tested Applications:

Cited Applications:

Human

Species Specificity:

Cited Species: human

Background Information

Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated α and β subunits (PMID: 9779984). CD11b, also known as Integrin alpha M or CR3A, belongs to the integrin alpha chain family. CD11b forms an α / β heterodimer with CD18 (integrin β 2). CD11b/CD18 is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complementcoated particles and pathogens (PMID: 9558116; 20008295). CD11b/CD18 is a receptor for the complement protein fragment iC3b, and is also a receptor for fibrinogen, factor X and ICAM1 (PMID: 2971974; 15485828).

Notable Publications

Author	Pubmed ID	Journal	Application
Cheng-Yu Liao	39025845	Nat Commun	FC
Huanrong Bai	38367493	Eur J Med Chem	FC


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1x10^6 human PBMCs were surface stained with 5 ul PE Anti-Human CD11b (PE-65116, Clone:ICRF 44), or PE Mouse IgG1 Isotype Control. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. Monocytes were gated.



1x10^6 human PBMCs were surface stained with 5 ul PE Anti-Human CD11b (PE-65116, Clone:ICRF 44), or PE Mouse IgG1 Isotype Control. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. Monocytes were gated.