## For Research Use Only

## PE Anti-Human CD8a (RPA-T8)

Catalog Number:PE-65144 1 Publications



**Basic Information** 

Catalog Number:

PE-65144

Size:

100tests, 5 μl/test

Source: Mouse

Isotype: IgG1, kappa

CD8a molecule Calculated MW: 235 aa, 26 kDa

UNIPROT ID:

P01732 Full Name:

BC025715

GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000153563

GenBank Accession Number:

**Purification Method:** Affinity purification

CloneNo.: RPA-T8

wavelengths:

Excitation/Emission maxima

496 nm, 565 nm / 578 nm

**Applications** 

**Tested Applications:** 

FC

**Cited Applications:** 

FC

Species Specificity:

human

**Background Information** 

 ${\tt CD8}\ is\ a\ transmembrane\ glycoprotein\ composed\ of\ two\ disulfide-linked\ chains.\ It\ can\ be\ present\ as\ a\ homodimer\ of\ present\ constant and\ constant\ constant$ CD8  $\alpha$  or as a heterodimer of CD8  $\alpha$  and CD8  $\beta$  (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8  $\alpha$   $\beta$  heterodimer while CD8  $\alpha$   $\alpha$  homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR)  $\gamma$   $\delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

**Notable Publications** 


Author	Pubmed ID	Journal	Application
Lin Yang	37106265	Clin Exp Med	FC

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA.

## Selected Validation Data



1x10^6 human PBMCs were surface stained with CL750 Anti-Human CD3 and 5 ul PE Anti-Human CD8a (PE-65144, Clone: RPA-T8) or PE Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



PE-65144 CD8a (RPA-T8)

1x10^6 human PBMCs were surface stained with 5 ul PE Anti-Human CD8a (PE-65144, Clone: PA-T8) (red) or PE Mouse IgG1 Isotype Control (blue). Cells were not fixed. Lymphocytes were gated.