For Research Use Only

PE Anti-Mouse CD8a (5H10-1)

Catalog Number: PE-65205



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Affinity purification

PE-65205

GeneID (NCBI):

BC030679

CloneNo.:

100ug, 0.2 mg/ml

12525

5H10-1

Source:

UNIPROT ID: P01731

Full Name:

Excitation/Emission maxima wavelengths:

Isotype: IgG2b, lambda

CD8 antigen, alpha chain

496 nm, 565 nm / 578 nm

Applications

Tested Applications:

FC

Species Specificity:

Mouse

Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α α homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) γ δ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Storage


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 0.5% BSA.

Selected Validation Data



1X10^6 C57BL/6 mouse splenocytes were surface co-stained with CoraLite® Plus 647 Anti-Mouse CD3 (17A2) and 0.1 ug PE Anti-Mouse CD8a (PE-65205, Clone: 5H10-1) or 0.1 ug PE Rat IgG2b Isotype Control (LTF-2). Cells were not fixed.