For Research Use Only

FcZero-rAb™ PE Anti-Human CD64 (10.1) Rabbit IgG Recombinant Antibody



Catalog Number: PE-FcA65572

Basic Information

Catalog Number:
PE-FcA65572
Concentration:

100tests, 5 ul/test Source: Rabbit Isotype:

gG Î

374 aa, 43 kDa

Purification Method: Protein A purification

Protein A purification CloneNo.: 10.1

Recommended Dilutions: FC: 5 ul per 10^6 cells in 100 $\,\mu$ l suspension

Excitation/Emission maxima

wavelengths: 496 nm, 565 nm / 578 nm

Applications

Tested Applications:

Species Specificity: human

Positive Controls:

FC: human PBMCs,

Background Information

Fc γ receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or inhibitory effects on cell functions after aggregation by complexed immunoglobulin G (IgG) (PMID: 17005690). CD64, also known as Fc γ RIA, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.

GenBank Accession Number:

Fc fragment of IgG, high affinity Ia,

BC032634

Full Name:

2209

GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000150337

receptor (CD64)

Calculated MW:


Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1x10^6 human PBMCs were surface stained with APC Anti-Human CD11b and 5 ul PE Anti-Human CD64 (10.1) Rabbit IgG Recombinant Antibody (PE-FcA65572, Clone:10.1) or PE Rabbit IgG Isotype Control Recombinant Antibody (PE-FcA98136, Clone: 240953C9). Cells were not fixed.