## For Research Use Only

## FcZero-rAb<sup>™</sup> PE Anti-Human VEcadherin/CD144 Rabbit Recombinant Antibody




Catalog Number:PE-FcA98071

Catalog Number: PE-FcA98071	GenBank Accession Number: NM_001795.5	Purification Method: Protein A purification	
Concentration: 100tests, 5 ul/test	GeneID (NCBI): 1003	CloneNo.: 240755B2	
Source: Rabbit Isotype: IgG	Full Name:Excitation/Emission maximacadherin 5, type 2 (vascularwavelengths:endothelium)496 nm, 565 nm / 578 nmCalculated MW:88 kDa	wavelengths:	
		Applications Tested Applications: FC	
Species Specificity: human			
play an important role in the cadherin), also known as Cadherin), also known as Cadherin), also known as Cadherin (PMID: 21: homophilic adhesion in the the organization of lateral endothelium (PMID: 152212: 18162609). The calculated m 140 kDa is higher due to pos	cadherin), also known as Cadherin-5 (CDH5) or CD144, is a member of the type II classical cadherin (V2- cadherin), also known as Cadherin-5 (CDH5) or CD144, is a member of the type II classical cadherin family of cell adhesion proteins (PMID: 21269602). VE-cadherin is expressed specifically in endothelial cells and mediates homophilic adhesion in the vascular endothelial (PMID: 1522121; 8555485; 21269602). VE-cadherin plays a role in the organization of lateral endothelial junctions and in the control of permeability properties of vascular endothelium (PMID: 1522121). VE-cadherin has also been shown to be required for angiogenesis (PMID: 16473763; 18162609). The calculated molecular weight of VE-cadherin is 88 kDa and the apparent molecular weight of 120- 140 kDa is higher due to post-translational glycosylation and phosphorylation (PMID: 10460833; 29894844). Full- length VE-cadherin can be proteolytically cleaved to generate a fragment of 90-100 kDa (PMID: 9786462;		
Storage: Store at 2-8°C. Avoid exposu Storage Buffer:	re to light. Stable for one year after shipme	ent.	
	PE-FcA98071 Concentration: 100tests, 5 ul/test Source: Rabbit Isotype: IgG Tested Applications: FC Species Specificity: human Cadherins are a family of tra play an important role in the cadherin), also known as Car adhesion proteins (PMID: 21: homophilic adhesion in the vi- the organization of lateral ei endothelium (PMID: 152212: 18162609). The calculated m 140 kDa is higher due to pos length VE-cadherin can be pi 22064597). Storage: Store at 2-8°C. Avoid exposu	PE-FcA98071 NM_001795.5   Concentration: GeneID (NCBI):   100tests, 5 ul/test 1003   Source: Full Name:   Rabbit cadherin 5, type 2 (vascular endothelium)   Isotype: endothelium)   IgG Calculated MW:   B8 kDa 88 kDa   On   Cadherins are a family of transmembrane glycoproteins that mediate of play an important role in the maintenance of normal tissue architecture cadherin), also known as Cadherin-5 (CDH5) or CD144, is a member of a dhesion proteins (PMID: 21269602). VE-cadherin is expressed specific homophilic adhesion in the vascular endothelium (PMID: 1522121; 855 the organization of lateral endothelial junctions and in the control of prendothelium (PMID: 1522121). VE-cadherin has also been shown to be a 18162609). The calculated molecular weight of VE-cadherin is 88 kDa a 140 kDa is higher due to post-translational glycosylation and phosphor length VE-cadherin can be proteolytically cleaved to generate a fragme 22064597).   Storage:   Storage:   Store at 2-8°C. Avoid exposure to light. Stable for one year after shipmed	

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10^6 HUVEC cells were surface stained with 5 ul PE Anti-Human VE-cadherin/CD144 Rabbit RecAb (PE-FcA98071, Clone:240755B2) (red) or PE Rabbit IgG Isotype Control RecAb (PE-FcA98136, Clone: 240953C9) (blue). Cells were not fixed.