For Research Use Only

PE-Cyanine7 Anti-Human PD-1/CD279 (EH12.2H7) Mouse IgG2a Recombinant Antibody




Catalog Number: PY7-65586


Basic Information	Catalog Number: PY7-65586	GenBank Accession Number: BC074740	Purification Method: Protein A purification
	Concentration: 100tests, 5 ul/test	GenelD (NCBI): 5133	CloneNo.: EH12.2H7
	Source: Mouse Isotype: IgG2a	Full Name: programmed cell death 1 Calculated MW: 288 aa, 32 kDa	Excitation/Emission maxima wavelengths: 450-500 nm / 778 nm
Species Specificity: human			
Background Information	Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).		
	hematopoietic cells in the pe PD-1 by its ligands PD-L1 or F cytolytic function (PMID: 194 Blockade of PD-1 can overcor	D-L2 transduces a signal that inhibits T-cel 26218). It is critical for the regulation of T	cytokines (PMID: 20636820). Engagement o l proliferation, cytokine production, and ell function during immunity and tolerance

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^6 PHA treated human PBMCs were surface stained with APC Anti-Human CD3 and 5 ul PE-Cyanine7 Anti-Human PD-1/CD279 (EH12.2H7) Mouse IgG2a Recombinant Antibody (PY7-65586, Clone:EH12.2H7), or 1x10^6 untreated human PBMCs were surface stained with APC Anti-Human CD3 and 5 ul PE-Cyanine7 Anti-Human PD-1/CD279 (EH12.2H7) Mouse IgG2a Recombinant Antibody (PY7-65586, Clone:EH12.2H7). Cells were not fixed.

1x10^6 PHA treated human PBMCs were surface stained with 5 ul PE-Cyanine7 Anti-Human PD-1/CD279 (EH12.2H7) Mouse IgG2a Recombinant Antibody (PY7-65586, Clone: EH12.2H7) (red) or unstained (blue). 1x10^6 untreated human PBMCs were surface stained with 5 ul PE-Cyanine7 Anti-Human PD-1/CD279 (EH12.2H7) Mouse IgG2a Recombinant Antibody (PY7-65586, Clone: EH12.2H7) (black, dashed). Cells were not fixed.