

For Research Use Only

FcZero-rAb® PE-Cyanine7 Anti-Mouse CD206 Rabbit Recombinant Antibody

Catalog Number:PY7-FcA98031



Basic Information

Catalog Number:

PY7-FcA98031

Concentration:

100ug, 200 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG0790

GenBank Accession Number:

NM_008625.2

GeneID (NCBI):

17533

UNIPROT ID:

Q2HZ94

Full Name:

mannose receptor, C type 1

Calculated MW:

165 kDa

Purification Method:

Protein A purification

CloneNo.:

240344D12

Recommended Dilutions:

FC: 0.1 ug per 10⁶ cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

496 nm, 565 nm / 778 nm

Applications

Tested Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC : mouse peritoneal macrophages,

Background Information

CD206, also named as MMR, CLEC13D and MRC1, is a type I membrane receptor that mediates the endocytosis of glycoproteins by macrophages. CD206 has been shown to bind high-mannose structures on the surface of potentially pathogenic viruses, bacteria, and fungi so that they can be neutralized by phagocytic engulfment. CD206 is a 170 kDa transmembrane protein which contains 5 domains: an amino-terminal cysteine-rich region, a fibronectin type II repeat, a series of eight tandem lectin-like carbohydrate recognition domains (responsible for the recognition of mannose and fucose), a transmembrane domain, and an intracellular carboxy-terminal tail. It is expressed on most tissue macrophages, in vitro derived dendritic cells, lymphatic and sinusoidal endothelia.

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ mouse peritoneal macrophages were surface stained with 0.1 ug PE-Cyanine7 Anti-Mouse CD206 Rabbit RecAb (PY7-FcA98031, Clone: 240344D12) or 0.1 ug FcZero-rAb® PE-Cyanine7 Rabbit IgG Isotype Control Recombinant Antibody (PY7-FcA98136, Clone: 240953C9). Cells were incubated with MonoZero™ Human/Mouse 3-in-1 blocking Reagent (PF00029) prior to staining. Cells were not fixed.



1x10⁶ mouse peritoneal macrophages were surface stained with 0.1 ug PE-Cyanine7 Anti-Mouse CD206 Rabbit RecAb (PY7-FcA98031, Clone: 240344D12) (red) or 0.1 ug FcZero-rAb® PE-Cyanine7 Rabbit IgG Isotype Control Recombinant Antibody (PY7-FcA98136, Clone: 240953C9) (blue). Cells were incubated with MonoZero™ Human/Mouse 3-in-1 blocking Reagent (PF00029) prior to staining. Cells were not fixed.