

For Research Use Only.
Not For Use In Diagnostics.

Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L)



Catalog Number:RGAM004

Information

Catalog Number:	RGAM004	Reactivity:	Mouse
Host:	Goat	Physical State:	Liquid
Applications:	IF, FC	Conjugation:	CoraLite® Plus 594

Recommended Dilutions

1:200-1:1000 for IF and FC

Fluorophore

CoraLite® Plus 594, Amax=588 nm, Emax=604 nm

Safety Notes

This product is for research use only, not for diagnostic or therapeutic use.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 50% glycerol, 10 mg/mL BSA, 0.1% Proclin-300, pH 7.4.
Aliquoting is unnecessary for -20°C storage

Purity


The antibody was purified from culture media supernatant by immunoaffinity chromatography using Protein G beads.

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunofluorescence of mouse cerebellum FFPE section was stained with Rabbit anti-NeuN polyclonal antibody (26975-1-AP, 1:200, green) and mouse anti-Calbindin-D28k monoclonal antibody (66394-1-Ig, 1:200, red). Multi-rAb CoraLite® Plus 488 conjugated Recombinant Goat anti-rabbit secondary antibody (RGAR002, 1:500) and Multi-rAb CoraLite® Plus 594 conjugated Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM004, 1:500) were used for detection.




Immunofluorescence of rat brain: rat brain FFPE section was stained with Rabbit anti-GFAP polyclonal antibody (16825-1-AP, 1:200, green) and mouse anti-NeuN monoclonal antibody (66836-1-Ig, red). Multi-rAb CoraLite® Plus 488 conjugated Recombinant Goat anti-rabbit secondary antibody (RGAR002, 1:500) and Multi-rAb CoraLite® Plus 594 conjugated Goat Anti-Mouse Recombinant Secondary Antibody (H+L) were used for detection (RGAM004, 1:500).




Immunofluorescence of MCF-7 cells: MCF-7 cells were fixed with 4% PFA and stained with Rabbit anti-ZO1 polyclonal antibody (21773-1-AP, 1:2000, green) and mouse anti-Alpha Tubulin monoclonal antibody (66031-1-Ig, 1:1000, red). Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002, 1:500) and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM004, 1:500) were used for detection.



Immunofluorescence of HeLa cells: HeLa cells were fixed with 4% PFA and stained with Rabbit anti-Alpha Tubulin polyclonal antibody (11224-1-AP, 1:200, green) and mouse anti-NPM1 monoclonal antibody (60096-1-Ig, 1:1000, red). Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002, 1:500) and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM004, 1:500) were used for detection.



Immunofluorescent analysis of (4% PFA) fixed OCT-embedded frozen mouse heart tissue using ACTC1-specific antibody (66125-1-Ig, Clone: 1F2B9) at dilution of 1:800 and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM004).



1X10⁶ MOLT4 were surface stained with 0.2 ug Anti-Human CD8 (65204-1-Ig, Clone: UCHT4) and Mouse IgG2a Isotype Control 66360-2-Ig. Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) RGAM004 was used at 1:500 for detection.