

For Research Use Only.  
Not For Use In Diagnostics.

# Multi-rAb™ Polymer HRP-Goat Anti-Rabbit/Mouse Universal Recombinant Secondary Antibody (H+L)



Catalog Number: RGAU011

## Information

### Catalog Number:

RGAU011

### Host:

Goat

### Applications:

IHC

### Reactivity:

Rabbit, Mouse

### Physical State:

Liquid

### Conjugation:

Polymer HRP

## Recommended Dilutions

Ready to use

## Safety Notes

This product is for research use only, not for diagnostic or therapeutic use.

## Storage

### Storage:

Store at 2-8°C. Stable for one year after shipment.

### Storage Buffer:

TBST + 10 mg/mL BSA + 1 mg/mL gelatin + 0.1% proclin300

Aliquoting is unnecessary for -20°C storage

## Purity

Protein G purification


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com) W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human skin cancer tissue slide using commercial IVD grade pan-CK mouse monoclonal antibody. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat Anti-Rabbit/Mouse Universal Recombinant Secondary Antibody (H+L) RGAU011 was used for detection.




Immunohistochemical analysis of paraffin-embedded human lymphoma tissue slide using other vendor's BCL2 rabbit recombinant antibody. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat Anti-Rabbit/Mouse Universal Recombinant Secondary Antibody (H+L) RGAU011 was used for detection.




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 80232-1-RR (Vimentin rabbit recombinant antibody) at dilution of 1:2500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-Rabbit/Mouse Universal Recombinant Secondary Antibody RGAU011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 66379-1-Ig (Beta catennin mouse monoclonal recombinant antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-Rabbit/Mouse Universal Recombinant Secondary Antibody RGAU011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 80001-1-RR (TDP-43 antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat Anti-Rabbit/Mouse Universal Recombinant secondary antibody RGAU011 was used for detection.




Immunohistochemical analysis of paraffin-embedded rat skeletal muscle tissue slide using 12715-1-AP (Dystrophin antibody) at dilution of 1:10000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat Anti-Rabbit/Mouse Universal Recombinant secondary antibody RGAU011 was used for detection.




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 81668-1-RR (F4/80 antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat Anti-Rabbit/Mouse Universal Recombinant secondary antibody RGAU011 was used for detection.




Immunohistochemical analysis of paraffin-embedded rat small intestine tissue slide using 66379-1-Ig (Beta Catenin antibody) at dilution of 1:20000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat Anti-Rabbit/Mouse Universal Recombinant secondary antibody RGAU011 was used for detection.




Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 66836-1-Ig (NeuN antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat Anti-Rabbit/Mouse Universal Recombinant secondary antibody RGAU011 was used for detection.




Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 66177-1-Ig (MPO antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat Anti-Rabbit/Mouse Universal Recombinant secondary antibody RGAU011 was used for detection.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 66786-1-Ig (SOX10 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat Anti-Rabbit/Mouse Universal Recombinant secondary antibody RGAU011 was used for detection.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 67300-1-Ig (CD107a / LAMP1 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat Anti-Rabbit/Mouse Universal Recombinant secondary antibody RGAU011 was used for detection.



Immunohistochemical analysis of paraffin-embedded rat kidney tissue slide using 67479-1-Ig (SLC22A7 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat Anti-Rabbit/Mouse Universal Recombinant secondary antibody RGAU011 was used for detection.