For Research Use Only

Multi-rAb™ IL-1 beta Multi-Recombinant antibody

Catalog Number:RMX00024




Basic Information	Catalog Number: RMX00024	GenBank Accession Number: BC008678	Purification Method: N/A				
	Concentration: 1000 µg/ml	GenelD (NCBI): 3553	Recommended Dilutions: WB 1:5000-1:50000				
	Source: Rabbit	ENSEMBL Gene ID: ENSG00000125538					
	lsotype: IgG	UNIPROT ID: P01584					
		Full Name: interleukin 1, beta Calculated MW: 269 aa, 31 kDa Observed MW: 30-35 kDa					
				Applications	Tested Applications: Positive Controls:		Controls:
					WB, ELISA Species Specificity: human	WB : LPS	and Brefeldin A treated THP-1 cells,
Background Information	Interleukin-1 is a pro-inflammatory cytokine with multiple biological effects. The IL-1 gene family encodes three proteins: IL-1 α , IL-1 β and their naturally occurring inhibitor IL-1RN. IL-1 Beta(IL-1 β), mainly produced by blood monocytes and tissue macrophages, has been implicated in mediating both acute and chronic inflammation. IL-1 β is known to be involved in a variety of cellular activities, including cell proliferation, differentiation and apoptosis. IL-1 β is emerging as a key mediator of carcinogenesis that characterizes host-environment interactions. Human IL-1 β is synthesized as a 31 kDa precursor. To gain activity, the precursor must be cleaved by caspase-1 between Asp116 and Ala117 to yield a 17 kDa mature form.						
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol. Aliquoting is unnecessary for -20°C storage						

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with RMX00024 (IL-1 beta antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.