

chromotek® anti-IgG Magnetic VHH Agarose for Immunoprecipitation

Product Code: mrlGma, mlGma, rIGma, mlG1ma, mlG2ama, mlG2bma, mlG3ma

Immunoprecipitation Protocol

Cell Material

The following protocol describes the preparation of a mammalian cell lysate.

For other types of cells, we recommend using 500 µg of cell extract.

Mammalian Cell Lysis

Note: Harvesting of cells and cell lysis should be performed with ice-cold buffers. We strongly recommend adding protease inhibitors to the lysis buffer to prevent degradation of your target protein and its binding partners.

For one immunoprecipitation reaction, we recommend using ~10⁶-10⁷ cells.

1. Choice of lysis buffer:
 - a. For cytoplasmic proteins, resuspend the cell pellet in 200 µL ice-cold lysis buffer by pipetting up and down. Supplement lysis buffer with protease inhibitor cocktail and 1 mM PMSF (not included).
 - b. For nuclear/chromatin proteins, resuspend cell pellet in 200 µL ice-cold RIPA buffer supplemented with DNase I (f.c. 75-150 Kunitz U/mL), MgCl₂ (f.c. 2.5 mM), protease inhibitor cocktail, and PMSF (f.c. 1 mM) (not included).
2. Place the tube on ice for 30 min and extensively pipette the suspension every 10 min.
3. Centrifuge cell lysate at 17,000x g for 10 min at +4°C. Transfer cleared lysate (supernatant) to a pre-cooled tube and add 300 µL dilution buffer supplemented with 1 mM PMSF and protease inhibitor cocktail (not included). If required, save 50 µL of diluted lysate for further analysis (input fraction).

Option 1: Incubation of lysate with primary antibody, followed by precipitation using beads:

Binding of protein of interest

1. Add 1 to 5 µg of primary rabbit or mouse antibody specific for protein of interest to the cleared, diluted lysate.
2. Rotate end-over-end for 1 hour at +4°C.

chromotek® anti-IgG Magnetic VHH Agarose for Immunoprecipitation

Product Code: mrlGma, mlGma, rlGma, mlG1ma, mlG2ama, mlG2bma, mlG3ma

Bead Equilibration

1. In the meantime, resuspend the beads by gently pipetting them up and down or by inverting the tube. Do not vortex the beads!
2. Transfer 25 μ L of bead slurry into a 1.5 mL reaction tube.
3. Add 500 μ L ice-cold dilution buffer.
4. Separate the beads with a magnet until the supernatant is clear. Discard the supernatant.

Immunoprecipitation

1. Add diluted lysate, including the primary antibody, to the equilibrated beads.
2. Rotate end-over-end for 1 hour at +4°C.

Option 2: Immobilization of primary antibody on beads followed by incubation with lysate:

Bead preparation:

1. Resuspend the beads by gently pipetting them up and down or by inverting the tube. Do not vortex the beads!
2. Transfer 25 μ L of bead slurry into a 1.5 mL reaction tube.
3. Add 500 μ L ice-cold dilution buffer.
4. Separate the beads with a magnet until the supernatant is clear. Discard the supernatant.
5. Add 500 μ L ice-cold dilution buffer.
6. Add 1 to 5 μ g of primary rabbit or mouse antibody specific for protein of interest.
7. Rotate end-over-end for 30 min at 4 °C.
8. Separate the beads with a magnet until the supernatant is clear. Discard the supernatant.

Immunoprecipitation

1. Add diluted lysate to the equilibrated beads comprising the primary antibody.
2. Rotate end-over-end for 1 hour at +4°C.

chromotek® anti-IgG Magnetic VHH Agarose for Immunoprecipitation

Product Code: mIgMma, mIgMma, rIgMma, mIg1ma, mIg2ama, mIg2bma, mIg3ma

Continue for all options:

Washing

1. Separate the beads with a magnet until the supernatant is clear.
2. If required, save 50 µL of supernatant for further analysis (flow-through/non-bound fraction).
3. Discard the remaining supernatant.
4. Resuspend beads in 500 µL wash buffer.
5. Separate the beads with a magnet until the supernatant is clear. Discard the remaining supernatant.
6. Repeat this step at least twice.
7. During the last washing step, transfer the beads to a new tube.

Optional: To increase stringency of the wash buffer, test various salt concentrations, e.g., 150-500 mM, and/or add a non-ionic detergent, e.g., Triton™ X-100 (see wash buffer compatibility information for maximum concentrations).

Elution with 2x SDS-sample buffer (Laemmli)

1. Remove the remaining supernatant.
2. Resuspend beads in 80 µL 2x SDS-sample buffer.
3. Boil beads for 5 min at +95°C to dissociate immunocomplexes from beads.
4. Separate the beads with a magnet.
5. Analyze the supernatant in SDS-PAGE / Western Blot.

Note: For Western blot detection, we recommend using a Western blot-validated antibody that is derived from a different species than the immunoprecipitation antibody or at least a different clone (if possible). Using different antibodies for immunoprecipitation and Western blot detection will (1) reassure you of the identity of the detected protein and (2) avoid the staining of the heavy and light chains of the immunoprecipitation antibody, which may mask the signal for your protein(s) of interest. For secondary antibodies for Western blot, we recommend Multi-rAb HRP-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (Proteintech RGAR001) or Multi-rAb™ HRP-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Proteintech RGAM001). If a Western blot detection antibody of a different species is unavailable, we recommend labeling the detection antibody using the FlexAble HRP Labeling kit (e.g., for rabbit IgG: Proteintech KFA005; for mouse IgG1: Proteintech KFA025), which will avoid staining of the heavy and light chains of the immunoprecipitation antibody.

chromotek[®] anti-IgG Magnetic VHH Agarose for Immunoprecipitation

Product Code: mrlGma, mlGma, rIGma, mlG1ma, mlG2ama, mlG2bma, mlG3ma

Related Products

| Product | Code |
|--|---|
| Anti-IgG VHH Agarose for Immunoprecipitation | mrlGa, mlGa, rIGa, mlG1a, mlG2aa, mlG2ba, mlG3a |

chromotek® anti-IgG Magnetic VHH Agarose for Immunoprecipitation

Product Code: mrlGma, mlGma, rlGma, mlG1ma, mlG2ama, mlG2bma, mlG3ma

Contact

Proteintech North America (HQ)

Proteintech Group, Inc.
5500 Pearl Street, Suite 400
Rosemont, IL 60018 USA

1-888-472-4522
proteintech@ptglab.com

Proteintech Europe

Transmission (6th Fl)
6 Atherton Street
M3 3GS, Manchester, UK

+44 161 839 3007
europe@ptglab.com

Proteintech Germany

Fraunhoferstr. 1
82152, Planegg-Martinsreid
Germany

+49 89 124 148 850
germany@ptglab.com

Disclaimer

For research use only (RUO), not for diagnostic or therapeutic use.

ChromoTek® is a registered trademark of ChromoTek GmbH, part of Proteintech Group.